

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2001 (21.06.2001)

PCT

(10) International Publication Number
WO 01/44452 A1

- (51) International Patent Classification⁷: **C12N 9/54**, (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (21) International Application Number: PCT/DK00/00660
- (22) International Filing Date: 1 December 2000 (01.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
PA 1999 01792 15 December 1999 (15.12.1999) DK
PA 2000 00708 1 May 2000 (01.05.2000) DK
PA 2000 01527 13 October 2000 (13.10.2000) DK
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant: **NOVOZYMES A/S** [DK/DK]; Krogshøjvej 36, DK-2880 Bagsvaerd (DK).
- Published:**
— With international search report.
- (72) Inventors: **FANØ, Tina, Sejersgaard**; Willemoesgade 17, 3. tv., DK-2100 København Ø (DK). **MIKKELSEN, Frank, F.**; Bykildevvej 5 st.tv., DK-2500 Valby (DK).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/44452 A1

(54) Title: SUBTILASE VARIANTS HAVING AN IMPROVED WASH PERFORMANCE ON EGG STAINS

(57) Abstract: The present invention relates to the use of a subtilase variant for removal of egg stains from laundry or from hard surfaces, where the subtilase variant comprises at least one additional amino acid residue in the active site loop (b) region from position 95 to 103 (BASBPN numbering). These subtilase variants are useful exhibiting excellent or improved wash performance on egg stains when used in e.g. cleaning or detergent compositions, including automatic dishwash compositions. The present invention also relates to novel subtilase variants, to isolated DNA sequences encoding the variants, expression vectors, host cells, and methods for producing and using the variants of the invention. Further, the present invention relates to cleaning and detergent compositions comprising the variants of the invention.

SUBTILASE VARIANTS HAVING AN IMPROVED WASH PERFORMANCE ON EGG STAINS

TECHNICAL FIELD

5 The present invention relates to the use of subtilase variants for removal of egg stains from laundry or from hard surfaces. In particular the present invention relates to the use of a subtilase variant for removal of egg stains from laundry or from hard surfaces, where the subtilase variant comprises at least
10 one additional amino acid residue in the active site loop (b) region from position 95 to 103 (BASBPN numbering, *vide infra*). These subtilase variants are useful exhibiting excellent or improved wash performance on egg stains when used in e.g. cleaning or detergent compositions, such as laundry detergent
15 compositions and dishwash composition, including automatic dishwash compositions. The present invention also relates to novel subtilase variants, to isolated DNA sequences encoding the variants, expression vectors, host cells, and methods for producing and using the variants of the invention. Further, the
20 present invention relates to cleaning and detergent compositions comprising the variants of the invention.

BACKGROUND OF THE INVENTION

25 In the detergent industry enzymes have for more than 30 years been implemented in washing formulations. Enzymes used in such formulations comprise proteases, lipases, amylases, cellulases, as well as other enzymes, or mixtures thereof. Commercially most important enzymes are proteases.

30 An increasing number of commercially used proteases are protein engineered variants of naturally occurring wild type proteases, e.g. DURAZYM[®] (Novo Nordisk A/S), RELEASE[®] (Novo Nordisk A/S),

MAXAPEM® (Gist-Brocades N.V.), PURAFECT® (Genencor International, Inc.).

Further, a number of protease variants are described in the art. A thorough list of prior art protease variants is given in WO
5 99/27082.

However, even though a number of useful protease variants have been described, there is still a need for new improved proteases or protease variants for a number of industrial uses.

10

In particular, the problem of removing egg stains from e.g. laundry or hard surfaces has been pronounced due to the fact that many proteases are inhibited by substances present in the egg white. Examples of such substances include trypsin inhibitor
15 type IV-0 (Ovo-inhibitor) and trypsin inhibitor type III-0 (Ovomucoid).

Therefore, an object of the present invention, is to provide improved subtilase variants, which are not, or which are only to
20 a limited extent, inhibited by such substances. A further object of the present invention is to provide improved subtilase variants, which are suitable for removal of egg stains from, for example, laundry and/or hard surfaces.

25 SUMMARY OF THE INVENTION

Thus, in a first aspect the present invention relates to the use of a subtilase variant for removal of egg stains from laundry or from hard surfaces, the subtilase variant comprising at least one additional amino acid residue in the active site loop (b) region from position 95 to 103
30 (BASBPN numbering).

In a second aspect the present invention relates to a subtilase variant selected from the group consisting of

5 a variant comprising at least one additional amino acid residue in the active site (b) loop corresponding to the insertion of at least one additional amino acid residue between positions 98 and 99 and further comprising at least one additional modification (BASBPN numbering), and

10 a variant comprising at least one additional amino acid residue in the active site (b) loop corresponding to the insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising at least one additional modification (BASBPN numbering),

15 where the variant - when tested in the "Ovo-inhibition Assay" disclosed in Example 4 herein - has a Residual Activity of at least 10%.

In a third aspect the present invention relates to a subtilase variant selected from the group consisting of

20

a variant comprising an insertion of at least one additional amino acid residue between positions 98 and 99 and further comprising a substitution in positions 133 and 143,

25 a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising a substitution in position 99,

30 a variant comprising an insertion of at least one additional amino acid residue between positions 98 and 99 and further comprising substitutions in positions 167, 170 and 194,

a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising an insertion of at least one additional amino acid residue between positions 216 and 217,

5

a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising an insertion of at least one additional amino acid residue between positions 217 and 218,

10

a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising an insertion of at least one additional amino acid residue between positions 42 and 43, and

15

a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising an insertion of at least one additional amino acid residue between positions 129 and 130.

20

In a fourth aspect the present invention relates to an isolated DNA sequence encoding a subtilase variant of the invention.

In a fifth aspect the present invention relates to an expression vector comprising the isolated DNA sequence of the invention.

25

In a sixth aspect the present invention relates to a microbial host cell transformed with the expression vector of the invention.

In a seventh aspect the present invention relates to a method for producing a subtilase variant according to the invention, wherein a host accord-

30

ing to the invention is cultured under conditions conducive to the expression and secretion of said variant, and the variant is recovered.

In an eight aspect the present invention relates to a cleaning
5 or detergent composition, preferably a laundry or dishwash
composition, comprising the variant of the invention.

In a ninth aspect the present invention relates to a method for
removal of egg stains from a hard surface or from laundry, the
10 method comprising contacting the egg stain-containing hard
surface or the egg stain-containing laundry with a cleaning or
detergent composition, preferably a laundry or dishwash
composition, containing a subtilase variant comprising at least
one additional amino acid residue in the active site loop (b)
15 region from position 95 to 103 (BASBPN numbering).

Still other aspect of the present invention will be apparent
from the below description and from the appended claims.

20 Concerning alignment and numbering reference is made to Fig. 1
which shows an alignments between subtilisin BPN' (a) (BASBPN)
and subtilisin 309 (BLSAVI) (b).

These alignments are in this patent application used as a
25 reference for numbering the residues.

DEFINITIONS

Prior to discussing this invention in further detail, the
following terms and conventions will first be defined.

30

NOMENCLATURE AND CONVENTIONS FOR DESIGNATION OF VARIANTS

In describing the various subtilase enzyme variants produced or contemplated according to the invention, the following nomenclatures and conventions have been adapted for ease of reference:

5

A frame of reference is first defined by aligning the isolated or parent enzyme with subtilisin BPN' (BASBPN).

The alignment can be obtained by the GAP routine of the GCG
10 package version 9.1 to number the variants using the following parameters: gap creation penalty = 8 and gap extension penalty = 8 and all other parameters kept at their default values.

Another method is to use known recognized alignments between
15 subtilases, such as the alignment indicated in WO 91/00345. In most cases the differences will not be of any importance.

Thereby a number of deletions and insertions will be defined in relation to BASBPN. In Fig. 1, subtilisin 309 (Savinase®) has 6
20 deletions in positions 36, 58, 158, 162, 163, and 164 in comparison to BASBPN. These deletions are in Fig. 1 indicated by asterixes (*).

The various modifications performed in a parent enzyme is
25 indicated in general using three elements as follows:

Original amino acid position substituted amino acid

The notation G195E thus means a substitution of a glycine in
30 position 195 with a glutamic acid.

In the case where the original amino acid residue may be any amino acid residue, a short hand notation may at times be used indicating only the position and substituted amino acid:

5 Position substituted amino acid

Such a notation is particular relevant in connection with modification(s) in homologous subtilases (*vide infra*).

- 10 Similarly when the identity of the substituting amino acid residue(s) is immaterial:

Original amino acid position

- 15 When both the original amino acid(s) and substituted amino acid(s) may comprise any amino acid, then only the position is indicated, e.g.: 170.

- When the original amino acid(s) and/or substituted amino acid(s) may comprise more than one, but not all amino acid(s), then the
20 selected amino acids are indicated inside brackets:

Original amino acid position {substituted amino acid₁, . . . , substituted amino acid_n}

- 25 For specific variants the specific three or one letter codes are used, including the codes Xaa and X to indicate any amino acid residue.

SUBSTITUTIONS:

- 30 The substitution of glutamic acid for glycine in position 195 is designated as:

Gly195Glu or G195E

or the substitution of any amino acid residue acid for glycine in position 195 is designated as:

5

Gly195Xaa or G195X

or

Gly195 or G195

10 The substitution of serine for any amino acid residue in position 170 would thus be designated

Xaa170Ser or X170S.

or

15 170Ser or 170S

Such a notation is particular relevant in connection with modification(s) in homologous subtilases (*vide infra*). 170Ser is thus meant to comprise e.g. both a Lys170Ser modification in
20 BASBPN and Arg170Ser modification in BLSAVI (cf. Fig. 1).

For a modification where the original amino acid(s) and/or substituted amino acid(s) may comprise more than one, but not all amino acid(s), the substitution of glycine, alanine, serine
25 or threonine for arginine in position 170 would be indicated by

Arg170{Gly,Ala,Ser,Thr} or R170{G,A,S,T}

to indicate the variants

30

R170G, R170A, R170S, and R170T.

DELETIONS:

A deletion of glycine in position 195 will be indicated by:

Gly195* or G195*

5

Correspondingly the deletion of more than one amino acid residue, such as the deletion of glycine and leucine in positions 195 and 196 will be designated

10

Gly195*+Leu196* or G195*+L196*

INSERTIONS:

The insertion of an additional amino acid residue such as e.g. a lysine after G195 is indicated by:

15

Gly195GlyLys or G195GK;

or, when more than one amino acid residue is inserted, such as e.g. a Lys, Ala and Ser after G195 this will be indicated as:

20

Gly195GlyLysAlaSer or G195GKAS (SEQ ID NO:1)

In such cases the inserted amino acid residue(s) are numbered by the addition of lower case letters to the position number of the amino acid residue preceding the inserted amino acid residue(s).

25 In the above example the sequences 194 to 196 would thus be:

194 195 196

BLSAVI A - G - L

194 195 195a 195b 195c 196

30

Variant A - G - K - A - S - L (SEQ ID NO:16)

In cases where an amino acid residue identical to the existing amino acid residue is inserted it is clear that a degeneracy in the nomenclature arises. If for example a glycine is inserted after the glycine in the above example this would be indicated
5 by G195GG. The same actual change could just as well be indicated as A194AG for the change from

194 195 196
BLSAVI A - G - L
10
to

194 195 195a 196
Variant A - G - G - L (SEQ ID NO:27)
15 194 194a 195 196

Such instances will be apparent to the skilled person, and the indication G195GG and corresponding indications for this type of insertions are thus meant to comprise such equivalent degenerate
20 indications.

FILLING A GAP:

Where a deletion in an enzyme exists in the reference comparison with the subtilisin BPN' sequence used for the numbering, an
25 insertion in such a position is indicated as:

*36Asp or *36D

for the insertion of an aspartic acid in position 36
30

MULTIPLE MODIFICATIONS:

Variants comprising multiple modifications are separated by pluses, e.g.:

Arg170Tyr+Gly195Glu or R170Y+G195E

5

representing modifications in positions 170 and 195 substituting tyrosine and glutamic acid for arginine and glycine, respectively.

10 Thus, Tyr167{Gly,Ala,Ser,Thr}+Arg170{Gly,Ala,Ser,Thr} designates the following variants:

	Tyr167Gly+Arg170Gly,	Tyr167Gly+Arg170Ala,
	Tyr167Gly+Arg170Ser,	Tyr167Gly+Arg170Thr,
15	Tyr167Ala+Arg170Gly,	Tyr167Ala+Arg170Ala,
	Tyr167Ala+Arg170Ser,	Tyr167Ala+Arg170Thr,
	Tyr167Ser+Arg170Gly,	Tyr167Ser+Arg170Ala,
	Tyr167Ser+Arg170Ser,	Tyr167Ser+Arg170Thr,
	Tyr167Thr+Arg170Gly,	Tyr167Thr+Arg170Ala,
20	Tyr167Thr+Arg170Ser,	and Tyr167Thr+Arg170Thr.

This nomenclature is particular relevant relating to modifications aimed at substituting, replacing, inserting or deleting amino acid residues having specific common properties, such as residues of positive charge (K, R, H), negative charge (D, E), or conservative amino acid modification(s) of e.g. Tyr167{Gly,Ala,Ser,Thr}+Arg170{Gly,Ala,Ser,Thr}, which signifies substituting a small amino acid for another small amino acid. See section "Detailed description of the invention" for further details.

30

Proteases

Enzymes cleaving the amide linkages in protein substrates are classified as proteases, or (interchangeably) peptidases (see Walsh, 1979, *Enzymatic Reaction Mechanisms*. W.H. Freeman and Company, San Francisco, Chapter 3).

5

Numbering of amino acid positions/residues

If nothing else is mentioned the amino acid numbering used herein correspond to that of the subtilase BPN' (BASBPN) sequence. For further description of the BPN' sequence, see Fig. 1 or Siezen et al., *Protein Engng.* 4 (1991) 719-737.

10

Serine proteases

A serine protease is an enzyme which catalyzes the hydrolysis of peptide bonds, and in which there is an essential serine residue at the active site (White, Handler and Smith, 1973 "*Principles of Biochemistry*," Fifth Edition, McGraw-Hill Book Company, NY, pp. 271-272).

15

The bacterial serine proteases have molecular weights in the 20,000 to 45,000 Dalton range. They are inhibited by diisopropylfluorophosphate. They hydrolyze simple terminal esters and are similar in activity to eukaryotic chymotrypsin, also a serine protease. A more narrow term, alkaline protease, covering a sub-group, reflects the high pH optimum of some of the serine proteases, from pH 9.0 to 11.0 (for review, see Priest (1977) *Bacteriological Rev.* 41 711-753).

20

25

Subtilases

A sub-group of the serine proteases tentatively designated subtilases has been proposed by Siezen et al., *Protein Engng.* 4 (1991) 719-737 and Siezen et al. *Protein Science* 6 (1997) 501-523. They are defined by homology analysis of more than 170

30

amino acid sequences of serine proteases previously referred to as subtilisin-like proteases. A subtilisin was previously often defined as a serine protease produced by Gram-positive bacteria or fungi, and according to Siezen et al. now is a subgroup of the subtilases. A wide variety of subtilases have been identified, and the amino acid sequence of a number of subtilases has been determined. For a more detailed description of such subtilases and their amino acid sequences reference is made to Siezen et al. (1997).

10

One subgroup of the subtilases, I-S1 or "true" subtilisins, comprises the "classical" subtilisins, such as subtilisin 168 (BSS168), subtilisin BPN' (SEQ ID NO:38), subtilisin Carlsberg (ALCALASE[®], NOVO NORDISK A/S), and subtilisin DY (BSSDY).

15

A further subgroup of the subtilases, I-S2 or high alkaline subtilisins, is recognized by Siezen et al. (*supra*). Sub-group I-S2 proteases are described as highly alkaline subtilisins and comprises enzymes such as subtilisin PB92 (BAALKP) (MAXACAL[®], Gist-Brocades NV), subtilisin 309 (SEQ ID NO:49) (SAVINASE[®], NOVO NORDISK A/S), subtilisin 147 (BLS147) (ESPERASE[®], NOVO NORDISK A/S), and alkaline elastase YaB (BSEYAB).

20

"SAVINASE[®]"

SAVINASE[®] is marketed by NOVO NORDISK A/S. It is subtilisin 309 from *B. Lentus* and differs from BAALKP only in one position (N87S, see Fig. 1 herein). SAVINASE[®] has the amino acid sequence designated b) in Fig. 1 and as shown in SEQ ID NO:49.

Parent subtilase

30

The term "parent subtilase" describes a subtilase defined according to Siezen et al. (1991 and 1997). For further details

see description of "SUBTILASES" immediately above. A parent subtilase may also be a subtilase isolated from a natural source, wherein subsequent modifications have been made while retaining the characteristic of a subtilase. Furthermore, a parent subtilase may also be a subtilase which has been prepared by the DNA shuffling technique, such as described by J.E. Ness et al., Nature Biotechnology, 17, 893-896 (1999). Alternatively the term "parent subtilase" may be termed "wild type subtilase".

10

Modification(s) of a subtilase variant

The term "modification(s)" used herein is defined to include chemical modification of a subtilase as well as genetic manipulation of the DNA encoding a subtilase. The modification(s) can be replacement(s) of the amino acid side chain(s), substitution(s), deletion(s) and/or insertions in or at the amino acid(s) of interest.

15

Subtilase variant

In the context of this invention, the term subtilase variant or mutated subtilase means a subtilase that has been produced by an organism which is expressing a mutant gene derived from a parent microorganism which possessed an original or parent gene and which produced a corresponding parent enzyme, the parent gene having been mutated in order to produce the mutant gene from which said mutated subtilase protease is produced when expressed in a suitable host.

20
25

Homologous subtilase sequences

Specific active site loop regions, and amino acid insertions in said loops of SAVINASE® subtilase are identified for modification herein to obtain a subtilase variant of the invention.

However, the invention is not limited to modifications of this particular subtilase, but extend to other parent (wild-type) subtilases, which have a homologous primary structure to that of SAVINASE®. The homology between two amino acid sequences is in this context described by the parameter "identity".

In order to determine the degree of identity between two subtilases the GAP routine of the GCG package version 9.1 can be applied (*infra*) using the same settings. The output from the routine is besides the amino acid alignment the calculation of the "Percent Identity" between the two sequences.

Based on this description it is routine for a person skilled in the art to identify suitable homologous subtilases and corresponding homologous active site loop regions, which can be modified according to the invention.

Isolated DNA sequence

The term "isolated", when applied to a DNA sequence molecule, denotes that the DNA sequence has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with

which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, 5 Dynan and Tijan, Nature 316:774-78, 1985). The term "an isolated DNA sequence" may alternatively be termed "a cloned DNA sequence".

Isolated protein

10 When applied to a protein, the term "isolated" indicates that the protein has been removed from its native environment.

In a preferred form, the isolated protein is substantially free of other proteins, particularly other homologous proteins (i.e. 15 "homologous impurities" (see below)).

An isolated protein is more than 10% pure, preferably more than 20% pure, more preferably more than 30% pure, as determined by SDS-PAGE. Further it is preferred to provide the protein in a 20 highly purified form, i.e., more than 40% pure, more than 60% pure, more than 80% pure, more preferably more than 95% pure, and most preferably more than 99% pure, as determined by SDS-PAGE.

25 The term "isolated protein" may alternatively be termed "purified protein".

Homologous impurities

The term "homologous impurities" means any impurity (e.g. another 30 polypeptide than the subtilase of the invention), which originate from the homologous cell where the subtilase of the invention is originally obtained from.

Obtained from

The term "obtained from" as used herein in connection with a specific microbial source, means that the polynucleotide and/or subtilase produced by the specific source, or by a cell in which a gene from the source has been inserted.

Substrate

The term "substrate" used in connection with a substrate for a protease should be interpreted in its broadest form as comprising a compound containing at least one peptide bond susceptible to hydrolysis by a subtilisin protease.

Product

The term "product" used in connection with a product derived from a protease enzymatic reaction should, in the context of the present invention, be interpreted to include the products of a hydrolysis reaction involving a subtilase protease. A product may be the substrate in a subsequent hydrolysis reaction.

20

Wash Performance

In the present context the term "wash performance" is used as an enzyme's ability to remove egg stains present on the object to be cleaned during e.g. wash or hard surface cleaning. See also the "Model Detergent Wash Performance Test" in Example 3 herein.

25

Performance Factor

The term "Performance Factor" is defined with respect to the below formula

30

$$P = R_{\text{variant}} - R_{\text{parent}}$$

wherein P is the Performance Factor, R_{variant} is the reflectance (measured at 460 nm) of the test material after being treated with a subtilase variant as described in the "Model Detergent Wash Performance Test", and R_{parent} is the reflectance (measured at 460 nm) of the test material after being treated with the corresponding parent subtilase as described in the "Model Detergent Wash Performance Test". For further details, see the "Model Detergent Wash Performance Test" in Example 3 herein.

10 Residual Activity

The term "Residual Activity" is defined as described in the "Ovo-inhibition Assay" herein (see Example 4).

BRIEF DESCRIPTION OF THE DRAWING

15 Fig. 1 shows an alignment between subtilisin BPN' (a) and Savinase[®](b) using the GAP routine mentioned above.

DETAILED DESCRIPTION OF THE INVENTION

The present inventors have found that subtilisin variants, wherein the active site loop (b) region is longer than those presently known, exhibit improved wash performance with respect to removal of egg stains. The identification thereof was done in constructing subtilisin variants, especially of the subtilisin 309 (BLSAVI or Savinase[®]), exhibiting improved wash performance properties (with respect to removal of egg stains) in model detergent compositions relative to the parent wild type enzyme.

Without being limited to any specific theory it is presently believed that the improved effect is due to an impeded binding of the egg white inhibitor in the active site loop (b) region of the subtilase variant. This in turn is probably due to structural changes of the active site loop (b) region because of

insertion of one or more additional amino acid residues in this particular site of the enzyme.

Thus, variants which are contemplated as being suitable for the
5 uses described herein are such variants where, when compared to the wild-type subtilase, one or more amino acid residues has been inserted in one or more of the following positions:
between positions 95 and 96, between positions 96 and 97, between positions 97 and 98, between positions 98 and 99, between positions 99 and
10 100, between positions 100 and 101, between positions 101 and 102, between positions 102 and 103, between positions 103 and 104, and combinations thereof.

Preferably, the insertion is made between position 97 and 98, between positions 98 and 99, between positions 99 and 100 and/or between positions
15 100 and 101, in particular between positions 98 and 99 and between positions 99 and 100.

A subtilase variant of the first aspect of the invention may be
20 a parent or wild-type subtilase identified and isolated from nature.

Such a parent wildtype subtilase may be specifically screened
for by standard techniques known in the art.

25

One preferred way of doing this may be by specifically PCR
amplify DNA regions known to encode active site loops in
subtilases from numerous different microorganism, preferably
different *Bacillus* strains.

30

Subtilases are a group of conserved enzymes, in the sense that their DNA and amino acid sequences are homologous. Accordingly it is possible to construct relatively specific primers flanking active site loops.

5

One way of doing this is by investigating an alignment of different subtilases (see e.g. Siezen et al. *Protein Science* 6 (1997) 501-523). It is from this routine work for a person skilled in the art to construct PCR primers flanking the active site loop corresponding to the active site loop (b) between amino acid residue 95 to 103 in any of the group I-S1 or I-S2 groups, such as from BLSAVI. Using such PCR primers to amplify DNA from a number of different microorganism, preferably different *Bacillus* strains, followed by DNA sequencing of said amplified PCR fragments, it will be possible to identify strains which produce subtilases of these groups comprising a longer, as compared to e.g. BLSAVI, active site region corresponding to the active site loop region from positions 95 to 103. Having identified the strain and a partial DNA sequence of such a subtilase of interest, it is routine work for a person skilled in the art to complete cloning, expression and purification of such a subtilase.

However, it is envisaged that a subtilase variant of the invention predominantly is a variant of a parent subtilase.

A subtilase variant suitable for the uses described herein, may be constructed by standard techniques known in the art such as by site-directed/random mutagenesis or by DNA shuffling of different subtilase sequences. See the "Material and Methods" section herein (*vide infra*) for further details.

As will be acknowledged by the skilled person, the variants described herein may, in addition to the at least one insertion from position 95 to 103, comprise at least one further modification. For example, the variants may comprise one or more
5 substitutions in the active site loop (b) region as well as one or more substitutions outside said region. Furthermore, the variants may comprise one or more further insertions outside the active site loop (b) region.

10 Moreover, the insertions in the regions described herein may encompass insertion of more than just one amino acid residue. For example the variant according to the invention may contain one insertion, two insertions, or more than two insertions, such as three, four or five insertions.

15 In preferred embodiments of the present invention, the further modification is performed in a position selected from the group consisting of: substitution in position 99, substitution in position 133, substitution in position 143, substitution in position 167, substitution
20 in position 170, substitution in position 194, insertion between positions 42 and 43, insertion between positions 129 and 130, insertion between positions 216 and 217, insertion between 217 and 218, and combinations thereof.

25 In an interesting embodiment of the invention the additional amino acid residue is inserted between position 98 and 99 (BASBPN numbering).

The insertion between position 98 and 99 is preferably selected
30 from the group consisting of (in BASBPN numbering)

X98X{A,T,G,S}, e.g., X98XA,X98XT,X98XG,X98XS;

X98X{D,E,K,R}, e.g., X98XD,X98XE,X98XK,X98XR;
X98X{H,V,C,N,Q}, e.g., X98XH,X98XV,X98XC,X98XN,X98XQ; and
X98X{F,I,L,M,P,W,Y}, e.g., X98XF,X98XI,X98XL,X98XM,X98XP,X98XW,
X98XY; preferably X98XA, X98XT, X98XG or X98XS;

5

or more specific for subtilisin 309 and closely related
subtilases, such as BAALKP, BLSUBL, and BSKSMK:

A98A{A,T,G,S}, e.g., A98AA,A98AT,A98AG,A98AS;
10 A98A{D,E,K,R}, e.g., A98AD,A98AE,A98AK,A98AR;
A98A{H,V,C,N,Q}, e.g., A98AH,A98AV,A98AC,A98AN,A98AQ;
A98A{F,I,L,M,P,W,Y}, e.g., A98AF,A98AI,A98AL,A98AM,A98AP,A98AW,
A98AY; preferably A98AA, A98AT, A98AG or A98AS.

15 Furthermore, it is presently preferred that the insertion
between position 98 and 99 is combined with a further
modification, namely substitution of an amino acid residue in
the positions 133 and 143, as well as substitution of an amino
acid residue in the positions 167, 170 and 194.

20

The substitutions (in addition to the insertion between position
98 and 99) in positions 133 and 134, respectively, are
preferably selected from the group consisting of

25 X133{A,T,G,S}, e.g., X133A,X133T,X133G,X133S;
X133{D,E,K,R}, e.g., X133D,X133E,X133K,X133R;
X133{H,V,C,N,Q}, e.g., X133H,X133V,X133C,X133N,X133Q;
X133{F,I,L,M,P,W,Y}, e.g., X133F,X133I,X133L,X133M,X133P,X133W,
X133Y;

30

X143{A,T,G,S}, e.g., X143A,X143T,X143G,X143S;
X143{D,E,K,R}, e.g., X143D,X143E,X143K,X143R;

X143{H,V,C,N,Q}, e.g., X143H,X143V,X143C,X143N,X143Q; and
X143{F,I,L,M,P,W,Y}, e.g., X143F,X143I,X143L,X143M,X143P,X143W,
X143Y.

- 5 In a preferred embodiment the substitution in position 133 is
selected from the group consisting of X133{D,E,K,R}, preferably
X133D or X133E, in particular X133E.

In another preferred embodiment the substitution in position 143
10 is selected from the group consisting of X143{D,E,K,R},
preferably X143K or X143R, in particular X143K.

An example of a preferred variant is a subtilase variant
comprising the following insertions and substitutions:

- 15 X98XS+X133E+X143K. A particular preferred variant is a savinase®
variant comprising the following insertions and substitutions:
A98AS+A133E+T143K.

Moreover, the substitutions (in addition to the insertion
20 between position 98 and 99) in positions 167, 170 and 134,
respectively, are preferably selected from the group consisting
of

- X167{A,T,G,S}, e.g., X167A,X167T,X167G,X167S;
25 X167{D,E,K,R}, e.g., X167D,X167E,X167K,X167R;
X167{H,V,C,N,Q}, e.g., X167H,X167V,X167C,X167N,X167Q;
X167{F,I,L,M,P,W,Y}, e.g., X167F,X167I,X167L,X167M,X167P,X167W,
X167Y;
30 X170{A,T,G,S}, e.g., X170A,X170T,X170G,X170S;
X170{D,E,K,R}, e.g., X170D,X170E,X170K,X170R;
X170{H,V,C,N,Q}, e.g., X170H,X170V,X170C,X170N,X170Q;

X170{F,I,L,M,P,W,Y}, e.g., X170F,X170I,X170L,X170M,X170P,X170W,
X170Y;

X194{A,T,G,S}, e.g., X194A,X194T,X194G,X194S;

5 X194{D,E,K,R}, e.g., X194D,X194E,X194K,X194R;

X194{H,V,C,N,Q}, e.g., X194H,X194V,X194C,X194N,X194Q; and

X194{F,I,L,M,P,W,Y}, e.g., X194F,X194I,X194L,X194M,X194P,X194W,
X194Y.

10 In a preferred embodiment the substitution in position 167 is
selected from the group consisting of X167{A,T,G,S}, in
particular X167A; the substitution in position 170 is selected
from the group consisting of X170{A,T,G,S}, in particular X170S;
and the substitution in position 194 is selected from the group
15 consisting of X194{F,I,L,M,P,W,Y}, in particular X194P.

An example of a preferred variant is a subtilase variant
comprising the following insertions and substitutions:

X98XT+X167A+X170S+X194P. A particular preferred variant is a
20 savinase® variant comprising the following insertions and
substitutions: A98AT+Y167A+R170S+A194P.

In a further interesting embodiment of the invention the
additional amino acid residue is inserted between position 99
25 and 100 (BASBPN numbering).

The insertion between position 99 and 100 is preferably selected
from the group consisting of (in BASBPN numbering)

30 X99X{A,T,G,S}, e.g., X99XA,X99XT,X99XG,X99XS;

X99X{D,E,K,R}, e.g., X99XD,X99XE,X99XK,X99XR;

X99X{H,V,C,N,Q}, e.g., X99XH,X99XV,X99XC,X99XN,X99XQ; and

X99X{F,I,L,M,P,W,Y}, e.g., X99XF,X99XI,X99XL,X99XM,X99XP,X99XW,
X99XY; preferably X99X{D,E,K,R}, in particular X99XD or X99XE;

or more specific for subtilisin 309 and closely related

5 subtilases, such as BAALKP, BLSUBL, and BSKSMK:

S99S{A,T,G,S}, e.g., S99SA,S99ST,S99SG,S99SS;

S99S{D,E,K,R}, e.g., S99SD,S99SE,S99SK,S99SR;

S99S{H,V,C,N,Q}, e.g., S99SH,S99SV,S99SC,S99SN,S99SQ;

10 S99S{F,I,L,M,P,W,Y}, e.g., S99SF,S99SI,S99SL,S99SM,S99SP,S99SW,
S99SY; preferably S99S{D,E,K,R}, in particular S99SD or S99SE.

With respect to insertions between position 99 and 100, it is -
in one interesting embodiment of the present invention -

15 preferred that the insertion is combined with a substitution in
position 99. Thus, in addition to the contemplated insertions
mentioned above, the following substitutions in position 99 are
considered relevant:

20 X99{A,T,G,S}, e.g., X99A,X99T,X99G,X99S;

X99{D,E,K,R}, e.g., X99D,X99E,X99K,X99R;

X99{H,V,C,N,Q}, e.g., X99H,X99V,X99C,X99N,X99Q; and

X99{F,I,L,M,P,W,Y}, e.g., X99F,X99I,X99L,X99M,X99P,X99W,X99Y.

25 In a preferred embodiment the substitution in position 99 is
selected from the group consisting of X99{A,T,G,S}, in
particular X99A or X99T.

An example of a preferred variant is a subtilase variant

30 comprising the following insertions and substitutions:

X99XD+X99A or X99XR+X99T. A particular preferred variant is a

savinase® variant comprising the following insertions and substitutions: S99SD+S99A or S99SR+S99T.

With respect to insertions between position 99 and 100, it is -
5 in another interesting embodiment of the present invention -
preferred that the insertion is combined with a further
insertion of at least one amino acid residue between positions
216 and 217. Thus, in addition to the contemplated insertions
mentioned above, the following insertions between positions 216
10 and 217 are considered relevant:

X216X{A,T,G,S}, e.g., X216XA,X216XT,X216XG,X216XS;
X216X{D,E,K,R}, e.g., X216XD,X216XE,X216XK,X216XR;
X216X{H,V,C,N,Q}, e.g., X216XH,X216XV,X216XC,X216XN,X216XQ; and
15 X216X{F,I,L,M,P,W,Y}, e.g., X216XF,X216XI,X216XL,X216XM,X216XP,
X216XW,X216XY.

In a preferred embodiment the insertion between positions 216
and 217 is selected from the group consisting of X216X{F,I,L,M,
20 P,W,Y} in particular X216XP.

Examples of preferred variants are subtilase variants comprising
the following insertions and substitutions: X99XD+X99A+X216XP as
well as X99XD+X99A+X216XDP. Particular preferred variants are
25 savinase® variants comprising the following insertions and
substitutions: S99SD+S99A+S216SP as well as S99SD+S99A+S216SDP.

With respect to insertions between position 99 and 100, it is -
in still another interesting embodiment of the present invention
30 - preferred that the insertion is combined with a further
insertion of at least one amino acid residue between positions
217 and 218. Thus, in addition to the contemplated insertions

mentioned above, the following insertions between positions 217 and 218 are considered relevant:

X217X{A,T,G,S}, e.g., X217XA,X217XT,X217XG,X217XS;
5 X217X{D,E,K,R}, e.g., X217XD,X217XE,X217XK,X217XR;
X217X{H,V,C,N,Q}, e.g., X217XH,X217XV,X217XC,X217XN,X217XQ; and
X217X{F,I,L,M,P,W,Y}, e.g., X217XF,X217XI,X217XL,X217XM,X217XP,
X217XW,X217XY.

10 In a preferred embodiment the insertion between positions 217 and 218 is selected from the group consisting of X217X{F,I,L,M,P,W,Y} in particular X217XP.

Examples of preferred variants are subtilase variants comprising
15 the following insertions and substitutions: X99XD+X99A+X217XP as well as X99XD+X217XP. Particular preferred variants are savinase® variants comprising the following insertions and substitutions: S99SD+S99A+L217LP as well as S99SD+L217P.

20 With respect to insertions between position 99 and 100, it is - in a further interesting embodiment of the present invention - preferred that the insertion is combined with a further insertion of at least one amino acid residue between positions 42 and 43. Thus, in addition to the contemplated insertions
25 mentioned above, the following insertions between positions 42 and 43 are considered relevant:

X42X{A,T,G,S}, e.g., X42XA,X42XT,X42XG,X42XS;
X42X{D,E,K,R}, e.g., X42XD,X42XE,X42XK,X42XR;
30 X42X{H,V,C,N,Q}, e.g., X42XH,X42XV,X42XC,X42XN,X42XQ; and
X42X{F,I,L,M,P,W,Y}, e.g., X42XF,X42XI,X42XL,X42XM,X42XP,
X42XW,X42XY.

In a preferred embodiment the insertion between positions 42 and 43 is selected from the group consisting of X42X{H,V,C,N,Q} in particular X42XN.

5

Examples of preferred variants are subtilase variants comprising the following insertions and substitutions: X99XD+X42XN as well as X99XD+X99A+X42XN. Particular preferred variants are savinase® variants comprising the following insertions and substitutions:
10 S99SD+D42DN as well as S99SD+S99A+D42DN.

With respect to insertions between position 99 and 100, it is -
in a still further interesting embodiment of the present
invention - preferred that the insertion is combined with a
15 further insertion of at least one amino acid residue between
positions 129 and 130. Thus, in addition to the contemplated
insertions mentioned above, the following insertions between
positions 129 and 130 are considered relevant:

20 X129X{A,T,G,S}, e.g., X129XA,X129XT,X129XG,X129XS;
X129X{D,E,K,R}, e.g., X129XD,X129XE,X129XK,X129XR;
X129X{H,V,C,N,Q}, e.g., X129XH,X129XV,X129XC,X129XN,X129XQ; and
X129X{F,I,L,M,P,W,Y}, e.g., X129XF,X129XI,X129XL,X129XM,X129XP,
X129XW,X129XY.

25

In a preferred embodiment the insertion between positions 129
and 130 is selected from the group consisting of X129X{D,E,K,R}.

Examples of preferred variants are subtilase variants comprising the following insertions and substitutions: X99XD+X129XD as well as X99XD+X99A+X129XD. Particular preferred variants are savinase® variants comprising the following insertions and
5 substitutions: S99SD+P129PD as well as S99SD+S99A+P129PD.

It is well known in the art that a so-called conservative substitution of one amino acid residue to a similar amino acid residue is expected to produce only a minor change in the
10 characteristic of the enzyme.

Table I below list groups of conservative amino acid substitutions.

Table I

<u>Conservative amino acid substitutions</u>	
<u>Common Property</u>	<u>Amino Acid</u>
Basic (positive charge)	K = lysine
	H = histidine
Acidic (negative charge)	E = glutamic acid
	D = aspartic acid
Polar	Q = glutamine
	N = asparagine
Hydrophobic	L = leucine
	I = isoleucine
	V = valine
	M = methionine
	F = phenylalanine
Aromatic	W = tryptophan
	Y = tyrosine
Small	G = glycine
	A = alanine
	S = serine
	T = threonine

According to this principle subtilase variants comprising
5 conservative substitutions are expected to exhibit
characteristics that are not drastically different from each
other.

Based on the disclosed and/or exemplified subtilase variants
10 herein, it is routine work for a person skilled in the art to
identify suitable conservative modification(s) to these variants
in order to obtain other subtilase variants exhibiting similarly
improved wash-performance.

It is preferred that the parent subtilase belongs to the subgroups I-S1 and I-S2, especially subgroup I-S2, both for isolating enzymes from nature or from the artificial creation of diversity, and for designing and producing variants from a parent subtilase.

In relation to variants from subgroup I-S1, it is preferred to select a parent subtilase from the group consisting of BSS168 (BSSAS, BSAPRJ, BSAPRN, BMSAMP), BASBPN, BSSDY, BLSCAR (BLKERA, BLSCA1, BLSCA2, BLSCA3), BSSPRC, and BSSPRD, or functional variants thereof having retained the characteristic of sub-group I-S1.

In relation to variants from subgroup I-S2 it is preferred to select a parent subtilase from the group consisting of BSAPRQ, BLS147 (BSAPRM, BAH101), BLSAVI (BSKSMK, BAALKP, BLSUBL), BYSYAB, BAPB92, TVTHER, and BSAPRS, or functional variants thereof having retained the characteristic of sub-group I-S2.

20

In particular, the parent subtilase is BLSAVI (Savinase®, NOVO NORDISK A/S), and a preferred subtilase variant of the invention is accordingly a variant of Savinase®. Thus, particular interesting variants are savinase® variants, i.e. BLSAVI variants,

25 wherein

1. Ser has been inserted between positions 98 and 99, Ala in position 133 has been substituted with Glu, and Thr in position 143 has been substituted with Lys (BASBPN numbering); or

30

2. Asp has been inserted between positions 99 and 100 and Ser in position 99 has been substituted with Ala (BASBPN numbering); or
3. Thr has been inserted between positions 98 and 99, Tyr in position 167 has been substituted with Ala, Arg in position 170 has been substituted with Ser, and Ala in position 194 has been substituted with Pro (BASBPN numbering); or
4. Asp has been inserted between positions 99 and 100, Ser in position 99 has been substituted with Ala, and Pro has been inserted between positions 217 and 218 (BASBPN numbering).
5. Asp has been inserted between positions 99 and 100, Ser in position 99 has been substituted with Ala, and Pro has been inserted between positions 216 and 217 (BASBPN numbering).
6. Asp has been inserted between positions 99 and 100, Ser in position 99 has been substituted with Ala, and Asp-Pro has been inserted between positions 216 and 217 (BASBPN numbering).
7. Asp has been inserted between positions 99 and 100, Ser in position 99 has been substituted with Ala, and Asp has been inserted between positions 129 and 130 (BASBPN numbering).
8. Asp has been inserted between positions 99 and 100, and Asn has been inserted between positions 42 and 43 (BASBPN numbering).
9. Asp has been inserted between positions 99 and 100, Ser in position 99 has been substituted with Ala, and Asn has been inserted between positions 42 and 43 (BASBPN numbering).

10. Arg has been inserted between positions 99 and 100, and Ser in position 99 has been substituted with Thr.

11. Asp has been inserted between positions 99 and 100, Ser in position 99 has been substituted with Ala, and Pro in
5 position 131 has been substituted with Thr.

The present invention also encompass use of any of the above mentioned subtilase variants in combination with any other modification to the amino acid sequence thereof. Especially combinations with other modifications known in the art to provide improved properties to the enzyme are envisaged. The art describes a number of subtilase variants with different improved properties and a number of those are mentioned in the "Background of the invention" section herein (*vide supra*). Those references are
10 disclosed here as references to identify a subtilase variant, which advantageously can be combined with a subtilase variant described herein.

Such combinations comprise the positions: 222 (improves oxidation stability), 218 (improves thermal stability), substitutions
20 in the Ca-binding sites stabilizing the enzyme, e.g. position 76, and many other apparent from the prior art.

In further embodiments a subtilase variant described herein may
25 advantageously be combined with one or more modification(s) in any of the positions:

27, 36, 56, 76, 87, 97, 101, 103, 104, 120, 123, 159, 167, 170, 206, 218, 222, 224, 232, 235, 236, 245, 248, 252 and 274.

30

Specifically the following BLSAVI, BLSUBL, BSKSMK, and BAALKP variants are considered appropriate for combination:

K27R, *36D, S56P, N76D, S87N, G97N, S101G, S103A, V104A, V104I,
V104N, V104Y, H120D, N123S, G159D, Y167, R170, Q206E, N218S,
M222S, M222A, T224S, A232V, K235L, Q236H, Q245R, N248D, N252K
5 and T274A.

Furthermore variants comprising any of the variants S101G+V104N,
S87N+S101G+V104N, K27R+V104Y+N123S+T274A, N76D+S103A+V104I or
N76D+V104A or other combinations of these mutations (V104N,
10 S101G, K27R, V104Y, N123S, T274A, N76D, V104A) or
S101G+S103A+V104I+G159D+A232V+Q236H+Q245R+N248D+N252K in combi-
nation with any one or more of the modification(s) mentioned
above exhibit improved properties.

15 Moreover, subtilase variants of the main aspect(s) of the inven-
tion are preferably combined with one or more modification(s) in
any of the positions 129, 131 and 194, preferably as 129K, 131H
and 194P modifications, and most preferably as P129K, P131H and
A194P modifications. Any of those modification(s) are expected
20 to provide a higher expression level of the subtilase variant in
the production thereof.

As mentioned above, the variants disclosed herein are only
inhibited by trypsin inhibitor type IV-0 to a limited extent
25 and, consequently, they exhibit excellent wash performance on
egg stains. Therefore, in order to enable the skilled person -
at an early stage of his development work - to select effective
and preferred variants for this purpose, the present inventors
have provided a suitable preliminary test, which can easily be
30 carried out by the skilled person in order to initially assess
the performance of the variant in question.

Thus, the "Ovo-inhibition Assay" disclosed in Example 4 herein may be employed to initially assess the potential of a selected variant. In other words, the "Ovo-inhibition Assay" may be employed to assess whether a selected variant will be inhibited, and to what extent, by the trypsin inhibitor type IV-0. Using this test, the suitability of a selected variant to remove egg stains can be assessed, the rationale being that if a selected variant is strongly inhibited by trypsin inhibitor type IV-0, it is normally not necessary to carry out further test experiments.

10

Therefore, a variant which is particularly interesting for the use described herein, is a variant which - when tested in the "Ovo-inhibition Assay" described in Example 4 herein - has a Residual Activity of at least 10%, e.g. at least 15%, such as at least 20%, preferably at least 25%, such as at least 30%, more preferably at least 35%. In a particular interesting embodiment of the invention, the variant has a Residual Activity of at least 40%, such as at least 45%, e.g. at least 50%, preferably at least 55%, such as at least 60%, more preferably at least 65%, such as at least 70%, even more preferably at least 75%, such as at least 80%, e.g. at least 90%, when tested in the "Ovo-inhibition Assay" described in Example 4 herein.

Evidently, it is preferred that the variant of the invention fulfils the above criteria on at least the stated lowest level, more preferably at the stated intermediate level and most preferably on the stated highest level.

Alternatively, or in addition to the above-mentioned assay, the suitability of a selected variant may be tested in the "Model Detergent Wash Performance Test" disclosed in Example 3 herein. The "Model Detergent Wash Performance Test" may be employed to

assess the ability of a variant, when incorporated in a standard detergent composition, to remove egg stains from a standard textile as compared to a reference system, namely the parent subtilase (incorporated in the same model detergent system and
5 tested under identical conditions). Using this test, the suitability of a selected variant to remove egg stains can be initially investigated, the rationale being that if a selected variant does not show a significant improvement in the test compared to the parent subtilase, it is normally not necessary
10 to carry out further test experiments.

Therefore, variants which are particular interesting for the use described herein, are such variants which, when tested in a model detergent composition comprising

15

	6.2%	LAS (Nansa 80S)
	2%	Sodium salt of C ₁₆ -C ₁₈ fatty acid
	4%	Non-ionic surfactant (Plurafax LF404)
	22%	Zeolite P
20	10.5%	Na ₂ CO ₃
	4%	Na ₂ Si ₂ O ₅
	2%	Carboxymethylcellulose (CMC)
	6.8%	Acrylate liquid CP5 40%
	20%	Sodium perborate (empirical formula NaBO ₂ .H ₂ O ₂)
25	0.2%	EDTA
	21%	Na ₂ SO ₄
		Water (balance)

as described in the "Model Detergent Wash Performance Test"
30 disclosed in Example 3 herein, shows an improved wash performance on egg stains as compared to the parent subtilase tested under identical conditions.

The improvement in the wash performance may be quantified by employing the so-called "Performance Factor" defined in Example 3, herein.

5

In a very interesting embodiment of the invention, the variant of the invention, when tested in the "Wash Performance Test" has a Performance Factor of at least 1, such as at least 1.5, e.g. at least 2, preferably at least 2.5, such as at least 3, e.g. at
10 least 3.5, in particular at least 4, such as at least 4.5, e.g. at least 5.

Evidently, it is preferred that the variant of the invention fulfils the above criteria on at least the stated lowest level,
15 more preferably at the stated intermediate level and most preferably on the stated highest level.

As indicated above, the present invention also provides novel subtilase variants. It will be understood that details and par-
20 ticulars concerning the novel subtilase variant aspects of the invention will be the same or analogous to the details and particulars of the variants discussed above in connection with the use aspect of the invention. This means that whenever appropriate, the statements concerning the use (e.g. preferred inser-
25 tions and substitutions, etc.) discussed in detail herein, apply mutatis mutandis to the novel subtilase variants according to the invention as well as to the method aspect and the cleaning and detergent composition aspect of the invention.

30 PRODUCING A SUBTILASE VARIANT

Many methods for cloning a subtilase and for introducing insertions into genes (e.g. subtilase genes) are well known in the

art, cf. the references cited in the "BACKGROUND OF THE INVENTION" section.

In general standard procedures for cloning of genes and introducing insertions (random and/or site directed) into said genes may be used in order to obtain a subtilase variant of the invention. For further description of suitable techniques reference is made to Examples herein (*vide infra*) and (Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus". John Wiley and Sons, 1990); and WO 96/34946.

Further, a subtilase variant may be constructed by standard techniques for artificial creation of diversity, such as by DNA shuffling of different subtilase genes (WO 95/22625; Stemmer WPC, Nature 370:389-91 (1994)). DNA shuffling of e.g. the gene encoding Savinase® with one or more partial subtilase sequences identified in nature to comprise an active site (b) loop regions longer than the active site (b) loop of Savinase®, will after subsequent screening for improved wash performance variants, provide subtilase variants suitable for the purposes described herein.

EXPRESSION VECTORS

A recombinant expression vector comprising a DNA construct encoding the enzyme of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures. The choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an

autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid.

5 Alternatively, the vector may be one that on introduction into a host cell is integrated into the host cell genome in part or in its entirety and replicated together with the chromosome(s) into which it has been integrated.

10 The vector is preferably an expression vector in which the DNA sequence encoding the enzyme of the invention is operably linked to additional segments required for transcription of the DNA. In general, the expression vector is derived from plasmid or viral DNA, or may contain elements of both. The term, "operably
15 linked" indicates that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in a promoter and proceeds through the DNA sequence coding for the enzyme.

20 The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell.

25 Examples of suitable promoters for use in bacterial host cells include the promoter of the *Bacillus stearothermophilus* maltogenic amylase gene, the *Bacillus licheniformis* alpha-amylase gene, the *Bacillus amyloliquefaciens* alpha-amylase gene, the *Bacillus subtilis* alkaline protease gen, or the *Bacillus*
30 *pumilus* xylosidase gene, or the phage Lambda P_R or P_L promoters or the *E. coli* lac, trp or tac promoters.

The DNA sequence encoding the enzyme of the invention may also, if necessary, be operably connected to a suitable terminator.

The recombinant vector of the invention may further comprise a
5 DNA sequence enabling the vector to replicate in the host cell in question.

The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, or a
10 gene encoding resistance to e.g. antibiotics like kanamycin, chloramphenicol, erythromycin, tetracycline, spectinomycin, or the like, or resistance to heavy metals or herbicides.

To direct an enzyme of the present invention into the secretory
15 pathway of the host cells, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) may be provided in the recombinant vector. The secretory signal sequence is joined to the DNA sequence encoding the enzyme in the correct reading frame. Secretory signal sequences are
20 commonly positioned 5' to the DNA sequence encoding the enzyme. The secretory signal sequence may be that normally associated with the enzyme or may be from a gene encoding another secreted protein.

25 The procedures used to ligate the DNA sequences coding for the present enzyme, the promoter and optionally the terminator and/or secretory signal sequence, respectively, or to assemble these sequences by suitable PCR amplification schemes, and to insert them into suitable vectors containing the information
30 necessary for replication or integration, are well known to persons skilled in the art (cf., for instance, Sambrook et al., op.cit.).

HOST CELL

The DNA sequence encoding the present enzyme introduced into the host cell may be either homologous or heterologous to the host in question. If homologous to the host cell, i.e. produced by the host cell in nature, it will typically be operably connected to another promoter sequence or, if applicable, another secretory signal sequence and/or terminator sequence than in its natural environment. The term "homologous" is intended to include a DNA sequence encoding an enzyme native to the host organism in question. The term "heterologous" is intended to include a DNA sequence not expressed by the host cell in nature. Thus, the DNA sequence may be from another organism, or it may be a synthetic sequence.

15

The host cell into which the DNA construct or the recombinant vector of the invention is introduced may be any cell which is capable of producing the present enzyme and includes bacteria, yeast, fungi and higher eukaryotic cells including plants.

20

Examples of bacterial host cells which, on cultivation, are capable of producing the enzyme of the invention are gram-positive bacteria such as strains of *Bacillus*, such as strains of *B. subtilis*, *B. licheniformis*, *B. lentus*, *B. brevis*, *B. stearothermophilus*, *B. alkalophilus*, *B. amyloliquefaciens*, *B. coagulans*, *B. circulans*, *B. lautus*, *B. megatherium* or *B. thuringiensis*, or strains of *Streptomyces*, such as *S. lividans* or *S. murinus*, or gram-negative bacteria such as *Echerichia coli*.

30

The transformation of the bacteria may be effected by protoplast transformation, electroporation, conjugation, or by using

competent cells in a manner known per se (cf. Sambrook et al., supra).

When expressing the enzyme in bacteria such as *E. coli*, the enzyme may be retained in the cytoplasm, typically as insoluble granules (known as inclusion bodies), or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed and the granules are recovered and denatured after which the enzyme is refolded by diluting the denaturing agent. In the latter case, the enzyme may be recovered from the periplasmic space by disrupting the cells, e.g. by sonication or osmotic shock, to release the contents of the periplasmic space and recovering the enzyme.

When expressing the enzyme in gram-positive bacteria such as *Bacillus* or *Streptomyces* strains, the enzyme may be retained in the cytoplasm, or may be directed to the extracellular medium by a bacterial secretion sequence. In the latter case, the enzyme may be recovered from the medium as described below.

METHOD FOR PRODUCING A SUBTILASE VARIANT

The present invention provides a method of producing an isolated enzyme according to the invention, wherein a suitable host cell, which has been transformed with a DNA sequence encoding the enzyme, is cultured under conditions permitting the production of the enzyme, and the resulting enzyme is recovered from the culture.

When an expression vector comprising a DNA sequence encoding the enzyme is transformed into a heterologous host cell it is possible to enable heterologous recombinant production of the enzyme of the invention.

Thereby it is possible to make a highly purified subtilase composition, characterized in being free from homologous impurities.

5

In this context homologous impurities means any impurities (e.g. other polypeptides than the enzyme of the invention) which originate from the homologous cell where the enzyme of the invention is originally obtained from.

10

The medium used to culture the transformed host cells may be any conventional medium suitable for growing the host cells in question. The expressed subtilase may conveniently be secreted into the culture medium and may be recovered therefrom by well-known procedures including separating the cells from the medium by centrifugation or filtration, precipitating proteinaceous components of the medium by means of a salt such as ammonium sulfate, followed by chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

20

CLEANING AND DETERGENT COMPOSITIONS

In general, cleaning and detergent compositions are well described in the art and reference is made to WO 96/34946; WO 97/07202; WO 95/30011 for further description of suitable cleaning and detergent compositions.

25

Furthermore the examples herein demonstrate the improvements in wash performance on egg stains for a number of subtilase variants.

30

Detergent Compositions

The subtilase variant may be added to and thus become a component of a detergent composition.

The detergent composition of the invention may for example be formulated as a hand or machine laundry detergent composition including a laundry additive composition suitable for pre-treatment of stained fabrics and a rinse added fabric softener composition, or be formulated as a detergent composition for use in general household hard surface cleaning operations, or be formulated for hand or machine dishwashing operations.

In a specific aspect, the invention provides a detergent additive comprising a subtilase enzyme of the invention. The detergent additive as well as the detergent composition may comprise one or more other enzymes such as another protease, a lipase, a cutinase, an amylase, a carbohydrase, a cellulase, a pectinase, a mannanase, an arabinase, a galactanase, a xylanase, an oxidase, e.g., a laccase, and/or a peroxidase.

In general the properties of the chosen enzyme(s) should be compatible with the selected detergent, (i.e. pH-optimum, compatibility with other enzymatic and non-enzymatic ingredients, etc.), and the enzyme(s) should be present in effective amounts.

Proteases: Suitable proteases include those of animal, vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. The protease may be a serine protease or a metallo protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from *Bacillus*, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and

subtilisin 168 (described in WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the *Fusarium* protease described in WO 89/06270 and WO 94/25583.

5

Examples of useful proteases are the variants described in WO 92/19729, WO 98/20115, WO 98/20116, and WO 98/34946, especially the variants with substitutions in one or more of the following positions: 27, 36, 57, 76, 87, 97, 101, 104, 120, 123, 167, 170,
10 194, 206, 218, 222, 224, 235 and 274.

Preferred commercially available protease enzymes include Alcalase™, Savinase™, Primase™, Duralase™, Esperase™, and Kannase™ (Novo Nordisk A/S), Maxatase™, Maxacal™, Maxapem™,
15 Properase™, Purafect™, Purafect OxP™, FN2™, and FN3™ (Genencor International Inc.).

Lipases: Suitable lipases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are
20 included. Examples of useful lipases include lipases from *Humicola* (synonym *Thermomyces*), e.g. from *H. lanuginosa* (*T. lanuginosus*) as described in EP 258 068 and EP 305 216 or from *H. insolens* as described in WO 96/13580, a *Pseudomonas* lipase, e.g. from *P. alcaligenes* or *P. pseudoalcaligenes* (EP 218 272),
25 *P. cepacia* (EP 331 376), *P. stutzeri* (GB 1,372,034), *P. fluorescens*, *Pseudomonas* sp. strain SD 705 (WO 95/06720 and WO 96/27002), *P. wisconsinensis* (WO 96/12012), a *Bacillus* lipase, e.g. from *B. subtilis* (Dartois et al. (1993), *Biochimica et Biophysica Acta*, 1131, 253-360), *B. stearothermophilus* (JP
30 64/744992) or *B. pumilus* (WO 91/16422).

Other examples are lipase variants such as those described in WO 92/05249, WO 94/01541, EP 407 225, EP 260 105, WO 95/35381, WO 96/00292, WO 95/30744, WO 94/25578, WO 95/14783, WO 95/22615, WO 97/04079 and WO 97/07202.

5

Preferred commercially available lipase enzymes include Lipolase™ and Lipolase Ultra™ (Novo Nordisk A/S).

Amylases: Suitable amylases (α and/or β) include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, α -amylases obtained from *Bacillus*, e.g. a special strain of *B. licheniformis*, described in more detail in GB 1,296,839.

15 Examples of useful amylases are the variants described in WO 94/02597, WO 94/18314, WO 96/23873, and WO 97/43424, especially the variants with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 181, 188, 190, 197, 202, 208, 209, 243, 264, 304, 305, 391, 408, and 444.

20

Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™ and BAN™ (Novo Nordisk A/S), Rapidase™ and Purastar™ (from Genencor International Inc.).

25 Cellulases: Suitable cellulases include those of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Suitable cellulases include cellulases from the genera *Bacillus*, *Pseudomonas*, *Humicola*, *Fusarium*, *Thielavia*, *Acremonium*, e.g. the fungal cellulases produced from *Humicola insolens*, *Myceliophthora thermophila* and *Fusarium oxysporum* disclosed in US 4,435,307, US 5,648,263, US 5,691,178, US 5,776,757 and WO 89/09259.

30

Especially suitable cellulases are the alkaline or neutral cellulases having colour care benefits. Examples of such cellulases are cellulases described in EP 0 495 257, EP 0 531 372, WO 96/11262, WO 96/29397, WO 98/08940. Other examples are cellulase variants such as those described in WO 94/07998, EP 0 531 315, US 5,457,046, US 5,686,593, US 5,763,254, WO 95/24471, WO 98/12307 and PCT/DK98/00299.

Commercially available cellulases include Celluzyme™, and Carezyme™ (Novo Nordisk A/S), Clazinase™, and Puradax HA™ (Genencor International Inc.), and KAC-500(B)™ (Kao Corporation).

Peroxidases/Oxidases: Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Examples of useful peroxidases include peroxidases from *Coprinus*, e.g. from *C. cinereus*, and variants thereof as those described in WO 93/24618, WO 95/10602, and WO 98/15257.

Commercially available peroxidases include Guardzyme™ (Novo Nordisk A/S).

The detergent enzyme(s) may be included in a detergent composition by adding separate additives containing one or more enzymes, or by adding a combined additive comprising all of these enzymes. A detergent additive of the invention, i.e. a separate additive or a combined additive, can be formulated e.g. as a granulate, a liquid, a slurry, etc. Preferred detergent additive formulations are granulates, in particular non-dusting granulates, liquids, in particular stabilized liquids, or slurries.

Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452 and may optionally be coated by methods known in the art. Examples of waxy coating materials are

5 poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molar weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty al-

10 cohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591. Liquid enzyme preparations may, for instance, be stabilized by adding a polyol such as propylene glycol, a sugar or sugar alco-

15 hol, lactic acid or boric acid according to established methods. Protected enzymes may be prepared according to the method disclosed in EP 238,216.

The detergent composition of the invention may be in any convenient form, e.g., a bar, a tablet, a powder, a granule, a paste

20 or a liquid. A liquid detergent may be aqueous, typically containing up to 70% water and 0-30% organic solvent, or non-aqueous.

25 The detergent composition typically comprises one or more surfactants, which may be non-ionic including semi-polar and/or anionic and/or cationic and/or zwitterionic. The surfactants are typically present at a level of from 0.1% to 60% by weight. When included therein the detergent will usually contain from

30 about 1% to about 40% of an anionic surfactant such as linear alkylbenzenesulfonate, alpha-olefinsulfonate, alkyl sulfate (fatty alcohol sulfate), alcohol ethoxysulfate, secondary al-

kanesulfonate, alpha-sulfo fatty acid methyl ester, alkyl- or alkenylsuccinic acid or soap.

When included therein the detergent will usually contain from
5 about 0.2% to about 40% of a non-ionic surfactant such as alcohol ethoxylate, nonylphenol ethoxylate, alkylpolyglycoside, alkyl dimethylamineoxide, ethoxylated fatty acid monoethanolamide, fatty acid monoethanolamide, polyhydroxy alkyl fatty acid amide, or N-acyl N-alkyl derivatives of glucosamine ("glucamides").

10

The detergent may contain 0-65% of a detergent builder or complexing agent such as zeolite, diphosphate, triphosphate, phosphonate, carbonate, citrate, nitrilotriacetic acid, ethylenediaminetetraacetic acid, diethylenetriaminepentaacetic acid, alkyl-
15 or alkenylsuccinic acid, soluble silicates or layered silicates (e.g. SKS-6 from Hoechst).

The detergent may comprise one or more polymers. Examples are carboxymethylcellulose, poly(vinylpyrrolidone), poly (ethylene
20 glycol), poly(vinyl alcohol), poly(vinylpyridine-N-oxide), poly(vinylimidazole), polycarboxylates such as polyacrylates, maleic/acrylic acid copolymers and lauryl methacrylate/acrylic acid copolymers.

25 The detergent may contain a bleaching system which may comprise a H_2O_2 source such as perborate or percarbonate which may be combined with a peracid-forming bleach activator such as tetraacetylethylenediamine or nonanoyloxybenzenesulfonate. Alternatively, the bleaching system may comprise peroxyacids of e.g.
30 the amide, imide, or sulfone type.

The enzyme(s) of the detergent composition of the invention may be stabilized using conventional stabilizing agents, e.g., a polyol such as propylene glycol or glycerol, a sugar or sugar alcohol, lactic acid, boric acid, or a boric acid derivative, e.g., an aromatic borate ester, or a phenyl boronic acid derivative such as 4-formylphenyl boronic acid, and the composition may be formulated as described in e.g. WO 92/19709 and WO 92/19708.

The detergent may also contain other conventional detergent ingredients such as e.g. fabric conditioners including clays, foam boosters, suds suppressors, anti-corrosion agents, soil-suspending agents, anti-soil redeposition agents, dyes, bactericides, optical brighteners, hydrotropes, tarnish inhibitors, or perfumes.

It is at present contemplated that in the detergent compositions any enzyme, in particular the enzyme of the invention, may be added in an amount corresponding to 0.01-100 mg of enzyme protein per liter of wash liquor, preferably 0.05-5 mg of enzyme protein per liter of wash liquor, in particular 0.1-1 mg of enzyme protein per liter of wash liquor.

The enzyme of the invention may additionally be incorporated in the detergent formulations disclosed in WO 97/07202 which is hereby incorporated as reference.

The invention is described in further detail in the following examples, which are not in any way intended to limit the scope of the invention as claimed.

In the detergent compositions, the abbreviated component identifications have the following meanings:

- 5 LAS: Sodium linear C₁₂ alkyl benzene sulphonate
- TAS: Sodium tallow alkyl sulphate
- XYAS: Sodium C_{1X} - C_{1Y} alkyl sulfate
- 10 SS: Secondary soap surfactant of formula 2-butyl octanoic acid
- 25EY: A C₁₂-C₁₅ predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide
- 15 45EY: A C₁₄-C₁₅ predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide
- 20 XYEZS: C_{1X}-C_{1Y} sodium alkyl sulfate condensed with an average of Z moles of ethylene oxide per mole
- Nonionic: C₁₃-C₁₅ mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5 sold under
- 25 the tradename Plurafax LF404 by BASF GmbH
- CFAA: C₁₂-C₁₄ alkyl N-methyl glucamide
- TFAA: C₁₆-C₁₈ alkyl N-methyl glucamide
- 30 Silicate: Amorphous Sodium Silicate (SiO₂:Na₂O ratio = 2.0)

- NaSKS-6: Crystalline layered silicate of formula $\delta\text{-Na}_2\text{Si}_2\text{O}_5$
- Carbonate: Anhydrous sodium carbonate
- 5 Phosphate: Sodium tripolyphosphate
- MA/AA: Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 80,000
- 10 Polyacrylate: Polyacrylate homopolymer with an average molecular weight of 8,000 sold under the trade-name PA30 by BASF GmbH
- Zeolite A: Hydrated Sodium Aluminosilicate of formula
15 $\text{Na}_{12}(\text{AlO}_2\text{SiO}_2)_{12} \cdot 27\text{H}_2\text{O}$ having a primary particle size in the range from 1 to 10 micrometers
- Citrate: Tri-sodium citrate dihydrate
- 20 Citric: Citric Acid
- Perborate: Anhydrous sodium perborate monohydrate bleach, empirical formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$
- 25 PB4: Anhydrous sodium perborate tetrahydrate
- Percarbonate: Anhydrous sodium percarbonate bleach of empirical formula $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$
- 30 TAED: Tetraacetyl ethylene diamine
- CMC: Sodium carboxymethyl cellulose

DETPMP: Diethylene triamine penta (methylene phosphonic acid), marketed by Monsanto under the Tradename Dequest 2060

5

PVP: Polyvinylpyrrolidone polymer

EDDS: Ethylenediamine-N, N'-disuccinic acid, [S,S] isomer in the form of the sodium salt

10

Suds 25% paraffin wax Mpt 50°C, 17% hydrophobic silica,
Suppressor: 58% paraffin oil

Granular Suds 12% Silicone/silica, 18% stearyl alcohol, 70%

15

suppressor: starch in granular form

Sulphate: Anhydrous sodium sulphate

20 HMWPEO: High molecular weight polyethylene oxide

TAE 25: Tallow alcohol ethoxylate (25)

Detergent Example I

25 A granular fabric cleaning composition in accordance with the invention may be prepared as follows:

30	Sodium linear C ₁₂ alkyl	6.5
	benzene sulfonate	
	Sodium sulfate	15.0
	Zeolite A	26.0

	Sodium nitrilotriacetate	5.0
	Enzyme	0.1
	PVP	0.5
	TAED	3.0
5	Boric acid	4.0
	Perborate	18.0
	Phenol sulphonate	0.1
	Minors	up to 100%

10 Detergent Example II

A compact granular fabric cleaning composition (density 800 g/l) in accord with the invention may be prepared as follows:

	45AS	8.0
15	25E3S	2.0
	25E5	3.0
	25E3	3.0
	TFAA	2.5
	Zeolite A	17.0
20	NaSKS-6	12.0
	Citric acid	3.0
	Carbonate	7.0
	MA/AA	5.0
	CMC	0.4
25	Enzyme	0.1
	TAED	6.0
	Percarbonate	22.0
	EDDS	0.3
	Granular suds suppressor	3.5
30	water/minors	Up to 100%

Detergent Example III

Granular fabric cleaning compositions in accordance with the invention which are especially useful in the laundering of coloured fabrics were prepared as follows:

5	LAS	10.7	-
	TAS	2.4	-
	TFAA	-	4.0
	45AS	3.1	10.0
	45E7	4.0	-
10	25E3S	-	3.0
	68E11	1.8	-
	25E5	-	8.0
	Citrate	15.0	7.0
	Carbonate	-	10.0
15	Citric acid	2.5	3.0
	Zeolite A	32.1	25.0
	Na-SKS-6	-	9.0
	MA/AA	5.0	5.0
	DETPMP	0.2	0.8
20	Enzyme	0.10	0.05
	Silicate	2.5	-
	Sulphate	5.2	3.0
	PVP	0.5	-
	Poly (4-vinylpyridine)-N-	-	0.2
25	Oxide/copolymer of vinyl- imidazole and vinyl- pyrrolidone		
	Perborate	1.0	-
30	Phenol sulfonate	0.2	-
	Water/Minors		Up to 100%

Detergent Example IV

Granular fabric cleaning compositions in accordance with the invention which provide "Softening through the wash" capability may be prepared as follows:

5			
	45AS	-	10.0
	LAS	7.6	-
	68AS	1.3	-
	45E7	4.0	-
10	25E3	-	5.0
	Coco-alkyl-dimethyl hydroxy-ethyl ammonium chloride	1.4	1.0
	Citrate	5.0	3.0
15	Na-SKS-6	-	11.0
	Zeolite A	15.0	15.0
	MA/AA	4.0	4.0
	DETPMP	0.4	0.4
	Perborate	15.0	-
20	Percarbonate	-	15.0
	TAED	5.0	5.0
	Smectite clay	10.0	10.0
	HMWPEO	-	0.1
	Enzyme	0.10	0.05
25	Silicate	3.0	5.0
	Carbonate	10.0	10.0
	Granular suds suppressor	1.0	4.0
	CMC	0.2	0.1
	Water/Minors	Up to 100%	

30

Detergent Example V

Heavy duty liquid fabric cleaning compositions in accordance with the invention may be prepared as follows:

5	LAS acid form	-	25.0
	Citric acid	5.0	2.0
	25AS acid form	8.0	-
	25AE2S acid form	3.0	-
	25AE7	8.0	-
10	CFAA	5	-
	DETPMP	1.0	1.0
	Fatty acid	8	-
	Oleic acid	-	1.0
	Ethanol	4.0	6.0
15	Propanediol	2.0	6.0
	Enzyme	0.10	0.05
	Coco-alkyl dimethyl hydroxy ethyl ammonium chloride	-	3.0
20	Smectite clay	-	5.0
	PVP	2.0	-
	Water / Minors	Up to 100%	

25 Powder automatic dishwash composition I

Nonionic surfactant	0.4	- 2.5%
Sodium metasilicate	0	- 20%
Sodium disilicate	3	- 20%
Sodium triphosphate	20	- 40%
Sodium carbonate	0	- 20%
Sodium perborate	2	- 9%

Tetraacetyl ethylene diamine (TAED)	1	- 4%
Sodium sulphate	5	- 33%
Enzymes	0.0001	- 0.1%

Powder automatic dishwash composition II

Nonionic surfactant (e.g. alcohol ethoxylate)	1	- 2%
Sodium disilicate	2	- 30%
Sodium carbonate	10	- 50%
Sodium phosphonate	0	- 5%
Trisodium citrate dihydrate	9	- 30%
Nitrilotrisodium acetate (NTA)	0	- 20%
Sodium perborate monohydrate	5	- 10%
Tetraacetyl ethylene diamine (TAED)	1	- 2%
Polyacrylate polymer (e.g. maleic acid/acrylic acid co-polymer)	6	- 25%
Enzymes	0.0001	- 0.1%
Perfume	0.1	- 0.5%
Water	5	- 10

5

Powder automatic dishwash composition III

Nonionic surfactant	0.5	- 2.0%
Sodium disilicate	25	- 40%
Sodium citrate	30	- 55%
Sodium carbonate	0	- 29%
Sodium bicarbonate	0	- 20%
Sodium perborate monohydrate	0	- 15%

Tetraacetyl ethylene diamine (TAED)	0	-	6%
Maleic acid/acrylic acid copolymer	0	-	5%
Clay	1	-	3%
Polyamino acids	0	-	20%
Sodium polyacrylate	0	-	8%
Enzymes	0.0001	-	0.1%

Powder automatic dishwash composition IV

Nonionic surfactant	1	-	2%
Zeolite MAP	15	-	42%
Sodium disilicate	30	-	34%
Sodium citrate	0	-	12%
Sodium carbonate	0	-	20%
Sodium perborate monohydrate	7	-	15%
Tetraacetyl ethylene diamine (TAED)	0	-	3%
Polymer	0	-	4%
Maleic acid/acrylic acid copolymer	0	-	5%
Organic phosphonate	0	-	4%
Clay	1	-	2%
Enzymes	0.0001	-	0.1%
Sodium sulphate	Balance		

5 Powder automatic dishwash composition V

Nonionic surfactant	1	-	7%
Sodium disilicate	18	-	30%
Trisodium citrate	10	-	24%
Sodium carbonate	12	-	20%

Monopersulphate (2 $\text{KHSO}_5 \cdot \text{KHSO}_4 \cdot \text{K}_2\text{SO}_4$)	15	- 21%
Bleach stabilizer	0.1	- 2%
Maleic acid/acrylic acid copolymer	0	- 6%
Diethylene triamine pentaacetate, pentasodium salt	0	- 2.5%
Enzymes	0.0001	- 0.1%
Sodium sulphate, water	Balance	

Powder and liquid dishwash composition with cleaning surfactant system VI

Nonionic surfactant	0	- 1.5%
Octadecyl dimethylamine N-oxide dihydrate	0	- 5%
80:20 wt.C18/C16 blend of octadecyl dimethylamine N-oxide dihydrate and hexadecyldimethyl amine N-oxide dihydrate	0	- 4%
70:30 wt.C18/C16 blend of octadecyl bis (hydroxyethyl)amine N-oxide anhydrous and hexadecyl bis (hydroxyethyl)amine N-oxide anhydrous	0	- 5%
C ₁₃ -C ₁₅ alkyl ethoxysulfate with an average degree of ethoxylation of 3	0	- 10%
C ₁₂ -C ₁₅ alkyl ethoxysulfate with an average degree of ethoxylation of 3	0	- 5%
C ₁₃ -C ₁₅ ethoxylated alcohol with an average degree of ethoxylation of 12	0	- 5%
A blend of C ₁₂ -C ₁₅ ethoxylated alcohols with an average degree of eth-	0	- 6.5%

oxylation of 9	
A blend of C ₁₃ -C ₁₅ ethoxylated alcohols with an average degree of ethoxylation of 30	0 - 4%
Sodium disilicate	0 - 33%
Sodium tripolyphosphate	0 - 46%
Sodium citrate	0 - 28%
Citric acid	0 - 29%
Sodium carbonate	0 - 20%
Sodium perborate monohydrate	0 - 11.5%
Tetraacetyl ethylene diamine (TAED)	0 - 4%
Maleic acid/acrylic acid copolymer	0 - 7.5%
Sodium sulphate	0 - 12.5%
Enzymes	0.0001 - 0.1%

Non-aqueous liquid automatic dishwashing composition VII

Liquid nonionic surfactant (e.g. alcohol ethoxylates)	2.0 - 10.0%
Alkali metal silicate	3.0 - 15.0%
Alkali metal phosphate	20.0 - 40.0%
Liquid carrier selected from higher glycols, polyglycols, polyoxides, glycolethers	25.0 - 45.0%
Stabilizer (e.g. a partial ester of phosphoric acid and a C ₁₆ -C ₁₈ alcohol)	0.5 - 7.0%
Foam suppressor (e.g. silicone)	0 - 1.5%
Enzymes	0.0001 - 0.1%

Non-aqueous liquid dishwashing composition VIII

Liquid nonionic surfactant (e.g. alcohol ethoxylates)	2.0	- 10.0%
Sodium silicate	3.0	- 15.0%
Alkali metal carbonate	7.0	- 20.0%
Sodium citrate	0.0	- 1.5%
Stabilizing system (e.g. mixtures of finely divided silicone and low molecular weight dialkyl polyglycol ethers)	0.5	- 7.0%
Low molecule weight polyacrylate polymer	5.0	- 15.0%
Clay gel thickener (e.g. bentonite)	0.0	- 10.0%
Hydroxypropyl cellulose polymer	0.0	- 0.6%
Enzymes	0.0001	- 0.1%
Liquid carrier selected from higher lycols, polyglycols, polyoxides and glycol ethers	Balance	

Thixotropic liquid automatic dishwashing composition IX

5

C ₁₂ -C ₁₄ fatty acid	0	- 0.5%
Block co-polymer surfactant	1.5	- 15.0%
Sodium citrate	0	- 12%
Sodium tripolyphosphate	0	- 15%
Sodium carbonate	0	- 8%
Aluminium tristearate	0	- 0.1%
Sodium cumene sulphonate	0	- 1.7%
Polyacrylate thickener	1.32	- 2.5%
Sodium polyacrylate	2.4	- 6.0%

Boric acid	0	- 4.0%
Sodium formate	0	- 0.45%
Calcium formate	0	- 0.2%
Sodium n-decyldiphenyl oxide disulphonate	0	- 4.0%
Monoethanol amine (MEA)	0	- 1.86%
Sodium hydroxide (50%)	1.9	- 9.3%
1,2-Propanediol	0	- 9.4%
Enzymes	0.0001	- 0.1%
Suds suppressor, dye, perfumes, water	Balance	

Liquid automatic dishwashing composition X

Alcohol ethoxylate	0	- 20%
Fatty acid ester sulphonate	0	- 30%
Sodium dodecyl sulphate	0	- 20%
Alkyl polyglycoside	0	- 21%
Oleic acid	0	- 10%
Sodium disilicate monohydrate	18	- 33%
Sodium citrate dihydrate	18	- 33%
Sodium stearate	0	- 2.5%
Sodium perborate monohydrate	0	- 13%
Tetraacetyl ethylene diamine (TAED)	0	- 8%
Maleic acid/acrylic acid copolymer	4	- 8%
Enzymes	0.0001	- 0.1%

5 Liquid automatic dishwashing composition containing protected bleach particles XI

Sodium silicate	5	- 10%
-----------------	---	-------

Tetrapotassium pyrophosphate	15	- 25%
Sodium triphosphate	0	- 2%
Potassium carbonate	4	- 8%
Protected bleach particles, e.g. chlorine	5	- 10%
Polymeric thickener	0.7	- 1.5%
Potassium hydroxide	0	- 2%
Enzymes	0.0001	- 0.1%
Water	Balance	

XII: Automatic dishwashing compositions as described in I, II, III, IV, VI and X, wherein perborate is replaced by per-
5 carbonate.

XIII: Automatic dishwashing compositions as described in I-VI, which additionally contain a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds described in "Effi-
10 cient manganese catalysts for low-temperature bleaching", Nature, (1994), 369, 637-639.

MATERIALS AND METHODS

15 TEXTILES:

WFK10N standard textile pieces (egg stains) were obtained from WFK Testgewebe GmbH, Christenfeld 10, D-41379 Brüggen-Bracht, Germany.

20 STRAINS:

B. subtilis DN1885 (Diderichsen et al., 1990).

B. lentus 309 and 147 are specific strains of *Bacillus lentus*, deposited with the NCIB and accorded the accession numbers NCIB 10309 and 10147, and described in US Patent No. 3,723,250 incorporated by reference herein.

5

E. coli MC 1000 (M.J. Casadaban and S.N. Cohen (1980); *J. Mol. Biol.* **138** 179-207), was made r^- , m^+ by conventional methods and is also described in US Patent Application Serial No. 039,298.

10 PLASMIDS:

pJS3 (SEQ ID NO:60): *E. coli* - *B. subtilis* shuttle vector containing a synthetic gene encoding for subtilase 309 (Described by Jacob Schiødt et al. in Protein and Peptide letters **3**:39-44 (1996)).

15

pSX222: *B. subtilis* expression vector (described in WO 96/34946).

GENERAL MOLECULAR BIOLOGY METHODS:

20 Unless otherwise mentioned the DNA manipulations and transformations were performed using standard methods of molecular biology (Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular
25 Biology". John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for *Bacillus*". John Wiley and Sons, 1990).

Enzymes for DNA manipulations were used according to the
30 specifications of the suppliers.

ENZYMES FOR DNA MANIPULATIONS

Unless otherwise mentioned all enzymes for DNA manipulations, such as e.g. restriction endonucleases, ligases etc., are obtained from New England Biolabs, Inc.

5 PROTEOLYTIC ACTIVITY

In the context of this invention proteolytic activity is expressed in Kilo NOVO Protease Units (KNPU). The activity is determined relatively to an enzyme standard (SAVINASE[®]), and the determination is based on the digestion of a dimethyl casein
10 (DMC) solution by the proteolytic enzyme at standard conditions, i.e. 50°C, pH 8.3, 9 min. reaction time, 3 min. measuring time. A folder AF 220/1 is available upon request to Novo Nordisk A/S, Denmark, which folder is hereby included by reference.

15 A GU is a Glycine Unit, defined as the proteolytic enzyme activity which, under standard conditions, during a 15 minutes' incubation at 40°C, with N-acetyl casein as substrate, produces an amount of NH₂-group equivalent to 1 mmole of glycine.

20 Enzyme activity can also be measured using the PNA assay, according to reaction with the soluble substrate succinyl-alanine-alanine-proline-phenyl-alanine-para-nitro-phenol, which is described in the Journal of American Oil Chemists Society, Rothgeb, T.M., Goodlander, B.D., Garrison, P.H., and Smith,
25 L.A., (1988).

FERMENTATION:

Fermentations for the production of subtilase enzymes were performed at 30°C on a rotary shaking table (300 r.p.m.) in 500
30 ml baffled Erlenmeyer flasks containing 100 ml BPX medium for 5 days.

Consequently in order to make an e.g. 2 liter broth 20
Erlenmeyer flasks were fermented simultaneously.

MEDIA:

5	BPX Medium Composition (per liter)	
	Potato starch	100 g
	Ground barley	50 g
	Soybean flour	20 g
	Na ₂ HPO ₄ × 12 H ₂ O	9 g
	Pluronic	0.1 g
	Sodium caseinate	10 g

The starch in the medium is liquefied with α -amylase and the
medium is sterilized by heating at 120°C for 45 minutes. After
sterilization the pH of the medium is adjusted to 9 by addition
10 of NaHCO₃ to 0.1 M.

EXAMPLE 1

CONSTRUCTION AND EXPRESSION OF ENZYME VARIANTS:

SITE-DIRECTED MUTAGENESIS:

15 Subtilase 309 (savinase®) site-directed variants of the
invention comprising specific insertions and comprising specific
substitutions were made by traditional cloning of DNA fragments
(Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd
Ed., Cold Spring Harbor, 1989) produced by PCR with oligos
20 containing the desired insertions (see below).

The template plasmid DNA was pJS3 (see below), or an analogue of
this containing a variant of Subtilase 309.

Insertions and substitutions were introduced by oligo directed mutagenesis to the construction of variants.

The Subtilase 309 variants were transformed into *E. coli*. DNA purified from a over night culture of these transformants were transformed into *B. subtilis* by restriction endonuclease digestion, purification of DNA fragments, ligation, transformation of *B. subtilis*. Transformation of *B. subtilis* was performed as described by Dubnau et al., 1971, J. Mol. Biol. 56, pp. 209-221.

10

SITE-DIRECTED MUTAGENESIS IN ORDER TO INTRODUCE INSERTIONS AND SUBSTITUTIONS IN A SPECIFIC REGION:

The overall strategy to used to perform site-directed mutagenesis was:

15 Mutagenic primers (oligonucleotides) were synthesized corresponding to the DNA sequence flanking the sites of insertion and substitutions, separated by the DNA base pairs defining the insertions and substitutions.

Subsequently, the resulting mutagenic primers were used in a PCR reaction with the modified plasmid pJS3 (see above). The resulting PCR fragment was purified and extended in a second PCR-reaction, the resulting PCR product was purified and either cloned into the *E. coli* - *B. subtilis* shuttle vector (see below) or extended in a third PCR-reaction before being digested by endonucleases and cloned into the *E. coli* - *B. subtilis* shuttle vector (see below). The PCR reactions are performed under normal conditions.

25

Following this strategy insertions and substitutions was constructed in savinase® wherein insertions and substitutions

was introduced according to the below table. The primers used for each PCR step are shown as well as the cloning sites used.

Following the above strategy a detailed example follows:

Two insertion and one substitution was constructed in savinase®
5 wherein insertions was introduced in position 99 (*99aD) and 217 (*217aP) respectively and a substitution was introduced in position S99A (see below).

The insertion and substitution at position 99 was introduced by a mutagenic primer (5' CCG AAC CTG AAC CAT CCG CGG CCC CTA GGA
10 CTT TAA CAG C 3' (sense)) (SEQ ID NO:71) were used in a PCR reaction with an opposite primer (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3' (antisense)) (SEQ ID NO:83).

The produced PCR fragment were extended towards the C-terminal of Savinase by a second round of PCR introducing the insertion
15 at position 217 with primer; 5' CAT CGA TGT ACC GTT TGG TAA GCT GGC ATA TGT TG 3' (SEQ ID NO:94). The second round PCR product were extended towards the C-terminal of Savinase by a third round of PCR with primer; 5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3' (SEQ ID NO:105), situated downstream at the Mlu I
20 site in pJS3. All PCR reactions used plasmid pJS3 as template. The extended DNA-fragment resulting from third round PCR was cloned into the Sal I- and Mlu I- sites of the modified plasmid pJS3 (see above).

The plasmid DNA was transformed into *E. coli* by well-known
25 techniques and one *E. coli* colony were sequenced to confirm the mutation designed.

All other variants were constructed in an analogous manner.

In order to purify a subtilase variant of the invention, the *B. subtilis* pJS3 expression plasmid comprising a variant of the

invention was transformed into a competent *B. subtilis* strain and was fermented as described above in a medium containing 10 µg/ml Chloramphenicol (CAM).

Primers and cloning sites:

Variant	Step 1 PCR primers	Step 2 PCR primers	Step 3 PCR	Cloning site
S99SR+S99T	Sense: (5' CAG AAG ATG TGG ACG CGC TTG 3') (SEQ ID NO:2) Antisense: (5' TGA ACC GCT GGT GGG GCC TAG GAC TTT AAC AG 3') (SEQ ID NO:7)	Sense: (step 1 PCR product) Antisense: (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:8)		HindIII-XbaI
S99SQ+S99T	Sense: (5' CAG AAG ATG TGG ACG CGC TTG 3') (SEQ ID NO:9) Antisense: (5' GAC CGA ACC TGA ACC CTG AGT GGC GCC TAG GAC 3') (SEQ ID NO:10)	Sense: (step 1 PCR product) Antisense: (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:11)		HindIII-XbaI
S99SD+M222S	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:12) Antisense: (5' GAC CGA ACC TGA ACC ATC GCT CGC CCC TAG GAC 3') (SEQ ID NO:13)	Sense: (step 1 PCR product) Antisense: (5' AGG AGT AGC CGA CGA TGT ACC GTT TAA GC 3') (SEQ ID NO:14)		SalI-MluI
L96LA+A98T+A108C+A138C	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:15) Antisense: (5' CCA TTC CAA TCC CTG GCA AAT CGA GCT GAC CGA ACC TGA ACC GCT GGT ACC CGC TAG GAC TTT AAC	Sense: (step 1 PCR product) Antisense: (5' AAC GCC TCT AGA AGT CGC GCT ATT AAC ACA TTG CTC GAG TGT GG 3') (SEQ ID NO:18)	Sense: (step 2 PCR product) Antisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:19)	SalI-MluI

	AGC G 3') (SEQ ID NO:17)			
A98AT+G97D	Sense: (5' CAG AAG ATG TGG ACG CGC TTG 3') (SEQ ID NO:20) Antisense: (5' AAC CGC TGG TGG CGT CTA GGA CTT TAA CAG CG 3') (SEQ ID NO:21)	Sense: (step1 PCR product) Antisense: (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:22)		HindIII- XbaI
A98AT+G97E	Sense: (5' CAG AAG ATG TGG ACG CGC TTG 3') (SEQ ID NO:23) Antisense: (5' AAC CGC TGG TGG CTT CTA GGA CTT TAA CAG CG 3') (SEQ ID NO:24)	Sense: (step 1 PCR prod- uct) Antisense: (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:25)		HindIII- XbaI
S99SA	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:26) Antisense: (5' ACC GAA CCT GAA CCT GCG CTC GCC CCT AGG 3') (SEQ ID NO:28)	Antisense: K828		HindIII- XbaI
S99SE+S99T	Sense: (5' CAG AAG ATG TGG ACG CGC TTG 3') (SEQ ID NO:29) Antisense: (5' GAC CGA ACC TGA GCC CTC GGT GGC GCC TAG GAC 3') (SEQ ID NO:30)	Antisense: (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:31)		HindIII- XbaI
S99SD+S99A+A133E	Sense: (5' CCC TTC GCC AAG TGA GAC TCT CGA GCA AGC TG 3') (SEQ ID NO:32) Antisense: (5' AAC CGC ACA GCG TTT	Sense: (5' AAA GTC CTA GGG GCC GCC GAC GGT TCA GGT TCG GTC AGC 3') (SEQ ID NO:34) Antisense: (step 1 PCR product)	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:35) Antisense: (step 2 PCR product)	SalI-MluI

	TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:33)			
S99SD+S99A+T143K	Sense: (5' TGT TAA TAG CGC GAA ATC CAG AGG CGT TCT TG 3') (SEQ ID NO:36) An- tisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:37)	Sense: (5' AAA GTC CTA GGG GCC GCC GAC GGT TCA GGT TCG GTC AGC 3') (SEQ ID NO:39) Antisense: (step 1 PCR product)	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:40) Antisense: (step 2 PCR product)	SalI-MluI
S99SD+S99A+S216SP	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:41) Antisense: (5' CCG AAC CTG AAC CAT CCG CGG CCC CTA GGA CTT TAA CAG C 3') (SEQ ID NO:42)	Sense: (step 1 PCR prod- uct) Antisense: (5' GAT GTA CCG TTT AAA GGG CTG GCA TAT GTT GAA CC 3') (SEQ ID NO:43)	Sense: (step 2 PCR product) Antisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:44)	SalI-MluI
S99SD+S99A+S216SDP	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3')SEQ ID NO:45) Antisense: (5' CCG AAC CTG AAC CAT CCG CGG CCC CTA GGA CTT TAA CAG C) SEQ ID NO:46)	Sense: (step 1 PCR prod- uct) Antisense: (5' GTA CCG TTT AAA GGA TCG CTG GCA TAT GTT GAA CC 3') (SEQ ID NO:47)	Sense: (step 2 PCR product) Antisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:48)	SalI-MluI
S99SD+S99A+P129PD	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:50) Antisense: (5' CCG AAC CTG AAC CAT CCG CGG CCC CTA GGA CTT TAA CAG C 3') (SEQ ID NO:51)	Sense: (step 1 PCR prod- uct) Antisense: (5' GTG TGG CAC TTG GCG AGT CAG GGC TTC CTA AAC TC 3') (SEQ ID NO:52)	Sense: (step 2 PCR product) Antisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:53)	SalI-MluI
S99SD+S99SA+P129PR	Sense: (5' GAG TTA AGC CCA GAA GAT	Antisense: (5' GTG TGG CACT TGG CGA	Sense: (step 2 PCR product)	SalI-MluI

	GTG GAC GCG 3') (SEQ ID NO:54) <i>Antisense:</i> (5' CCG AAC CTG AAC CAT CCG CGG CCC CTA GGA CTT TAA CAG C 3') (SEQ ID NO:55)	TCG AGG GCT TCC TAA ACT C 3') (SEQ ID NO:56)	<i>Antisense:</i> (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:57)	
S99SD+S99A+L217F+ A228V+A230V	<i>Sense:</i> (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:58) <i>Antisense:</i> (5' CCG AAC CTG AAC CAT CCG CGG CCC CTA GGA CTT TAA CAG C 3') (SEQ ID NO:59)	<i>Anstisense:</i> (5' (AAG GGC GGC CAC ACC TAC AAC ATG AGG AGT AGC CAT CGA TGT ACC GTT AAA GCT GGC ATA TGT TGA AC 5') (SEQ ID NO:61)	<i>Sense:</i> (step 2 PCR product) <i>Antisense:</i> (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:62)	SalI-MluI
S99SD+S99A+L217LP	<i>Sense:</i> (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:63) <i>Antisense:</i> (5' CCG AAC CTG AAC CAT CCG CGG CCC CTA GGA CTT TAA CAG C 3') (SEQ ID NO:64)	<i>Antisense:</i> (5' CAT CGA TGT ACC GTT TGG TAA GCT GGC ATA TGT TG 3') (SEQ ID NO:65)	<i>Sense:</i> (step 2 PCR product) <i>Antisense:</i> (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:66)	SalI-MluI

Variant	Step 1 PCR primers	Step 2 PCR primers	Step 3 PCR	Cloning site
G97GI+S99T	<p><i>Sense:</i> (5' GCT GTT AAA GTC CTA GGG ATC GCG ACT GGT TCA GGT TCG GTC AGC 3') (SEQ ID NO:76)</p> <p><i>Antisense:</i> (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:68)</p>			AvrII-XbaI
A98AS+A133E+T143K	<p><i>Sense:</i> (5' GTT AAA GTC CTA GGG GCG TCG AGC GGT TCA GGT TCG GTC 3') (SEQ ID NO:69)</p> <p><i>Antisense:</i> (5' C AAG AAC GCC TCT AGA TTT CGC GCT ATT AAC AGC TTG CTC GAG TGT TTC ACT TGG CGA AGG GCT TCC 3') (SEQ ID NO:70)</p>			AvrII-XbaI
A98AG	<p><i>Sense:</i> (5' GCT GTT AAA GTC CTA GGG GCG GGT AGC GGT TCA GGT TCG GTC 3') (SEQ ID NO:72)</p> <p><i>Antisense:</i> (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:73)</p>			AvrII-XbaI
A98AS+R45K+S105G	<p><i>Sense:</i> (5' GTC CTC GAT ACA GGG ATA TCC ACT CAT CCA GAT CTA AAT ATT AAA GGT GGC GCA AGC TTT GTA C 3') (SEQ ID NO:74)</p> <p><i>Antisense:</i> (5' CGC CCC TAG GAC TTT AAC AGC 3') (SEQ ID NO:75)</p>	<p><i>Sense:</i> (5' GCT GTT AAA GTC CTA GGG GCG TCG AGC GGT TCA GGT TCG GTC GGG TCG ATT GCC CAA GGA TTG 3') (SEQ ID NO:76)</p> <p><i>Antisense:</i> (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:77)</p>	<p><i>Sense:</i> (step 1 PCR product)</p> <p><i>Antisense:</i> (step 2 PCR product)</p>	EcoRV-XbaI

A98ASGTG	<p><i>Sense:</i> (5' GCT GTT AAA GTC CTA GGG GCG TCG GGC ACT GGC AGC GGT TCA GGT TCG GTC 3') (SEQ ID NO:78) <i>Antisense:</i> (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:79)</p>			AvrII-XbaI
A98AP+A98G+S99A	<p><i>Sense:</i> (5' GCT GTT AAA GTC CTA GGG GGC CCA GCC GGT TCA GGT TCG GTC AGC 3') (SEQ ID NO:80) <i>Antisense:</i> (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:81)</p>			AvrII-XbaI
A98AI+A98G+S99H+G100S+S101A	<p><i>Sense:</i> (5' GCT GTT AAA GTC CTA GGG GGC ATC CAT TCG GCA GGT TCG GTC AGC TCG ATT 3') (SEQ ID NO:82) <i>Antisense:</i> (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:84)</p>			AvrII-XbaI
S99SD+S99A	<p><i>Sense:</i> (5' GCT GTT AAA GTC CTA GGG GCG GCA GAC GGT TCA GGT TCG GTC AGC 3') (SEQ ID NO:85) <i>Antisense:</i> (5' GAT TAA CGC GTT GCC GCT TCT G 3') (SEQ ID NO:86)</p>			AvrII-XbaI
S99SD+S99A+P131T	<p><i>Sense:</i> (5' GCT GTT AAA GTC CTA GGG GCG GCA GAC GGT TCA GGT TCG GTC AGC</p>			AvrII-XbaI

	TCG ATT GCC CAA GGA TTG 3') (SEQ ID NO:87) Antisense: (5' TTG CTC GAG TGT GGC ACT GGT CGA AGG GCT TCC TAA ACT 3') (SEQ ID NO:88)			
Variant	Step 1 PCR primers	Step 2 PCR primers	Step 3 PCR	Cloning site
L96LA	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:89) Antisense: (5' AAC CGC TCG CCC CTG CTA GGA CTT TAA CAG 3') (SEQ ID NO:90)	Sense: (step 1 PCR prod- uct) Anstisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TG C 3') (SEQ ID NO:91)		SalI-MluI
S99SN	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:92) Antisense: (5' GAC CGA ACC TGA ACC GTT GCT CGC CCC TAG GAC 3') (SEQ ID NO:93)	Sense: (step 1 PCR prod- uct) Anstisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TG C 3') (SEQ ID NO:95)		SalI-MluI
S99SD	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:96) Antisense: (5' GAC CGA ACC TGA ACC ATC GCT CGC CCC TAG GAC 3') (SEQ ID NO:97)	Sense: (step 1 PCR prod- uct) Anstisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TG C 3') (SEQ ID NO:98)		SalI-MluI
S99SE	Sense: (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:99) Antisense: (5' GAC CGA ACC TGA ACC TTC GCT CGC CCC TAG GAC 3') (SEQ ID	Sense: (step 1 PCR prod- uct) Anstisense: (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TG C 3') (SEQ ID NO:101)		SalI-MluI

	NO:100)			
A98AT+Y167A+R170S+A194P	<p><i>Sense:</i> (5' GAG TTA AGC CCA GAA GAT GTG GAC GCG 3') (SEQ ID NO:102)</p> <p><i>Antisense:</i> (5' TGT GTA AAG TAA CTC ATT TGG TGA GCC AG 3') (SEQ ID NO:103)</p>	<p><i>Sense:</i> (step 1 PCR product)</p> <p><i>Anstisense:</i> (5' CCG ACT GCC ATT GCG TTC GCA TAC GAC GCC GGG GCG CTG ATT GAG CCT GCA C 3') (SEQ ID NO:104)</p>	<p><i>Sense:</i> (step 2 PCR product)</p> <p><i>Anstisense:</i> (5' CTG CAC GTT TAC CCC GGG TGC GAC AAT GTC AAG GCC TGG GCC ATA CTG TG 3') (SEQ ID NO:3)</p>	SallI-XmaI
S99SD+D42DN+S99A	<p><i>Sense:</i> (5' CTC GAT ACA GGG ATA TCC ACT CAT CCA GAT CTA AAC AAT ATT CGT GGT GGC G 3') (SEQ ID NO:4)</p> <p><i>Antisense:</i> (5' CCG AAC CTG AAC CAT CCG CGG CCC CTA GGA CTT TAA CAG C 3') (SEQ ID NO:5)</p>	<p><i>Sense:</i> (step 1 PCR product)</p> <p><i>Anstisense:</i> (5' AAC CGC ACA GCG TTT TTT TAT TGA TTA ACG CGT TGC 3') (SEQ ID NO:6)</p>		EcoRV-MluI

EXAMPLE 2**PURIFICATION OF ENZYME VARIANTS:**

This procedure relates to purification of a 2 liter scale
5 fermentation for the production of the subtilases of the
invention in a *Bacillus* host cell.

Approximately 1.6 liters of fermentation broth were centrifuged
at 5000 rpm for 35 minutes in 1 liter beakers. The supernatants
10 were adjusted to pH 6.5 using 10% acetic acid and filtered on
Seitz Supra S100 filter plates.

The filtrates were concentrated to approximately 400 ml using an
Amicon CH2A UF unit equipped with an Amicon S1Y10 UF cartridge.
15 The UF concentrate was centrifuged and filtered prior to
absorption at room temperature on a Bacitracin affinity column
at pH 7. The protease was eluted from the Bacitracin column at

room temperature using 25% 2-propanol and 1 M sodium chloride in a buffer solution with 0.01 dimethylglutaric acid, 0.1 M boric acid and 0.002 M calcium chloride adjusted to pH 7.

- 5 The fractions with protease activity from the Bacitracin purification step were combined and applied to a 750 ml Sephadex G25 column (5 cm dia.) equilibrated with a buffer containing 0.01 dimethylglutaric acid, 0.2 M boric acid and 0.002 m calcium chloride adjusted to pH 6.5.

10

Fractions with proteolytic activity from the Sephadex G25 column were combined and applied to a 150 ml CM Sepharose CL 6B cation exchange column (5 cm dia.) equilibrated with a buffer containing 0.01 M dimethylglutaric acid, 0.2 M boric acid, and
15 0.002 M calcium chloride adjusted to pH 6.5.

The protease was eluted using a linear gradient of 0-0.1 M sodium chloride in 2 litres of the same buffer (0-0.2 M sodium chloride in case of Subtilisin 147).

20

In a final purification step protease containing fractions from the CM Sepharose column were combined and concentrated in an Amicon ultrafiltration cell equipped with a GR81PP membrane (from the Danish Sugar Factories Inc.).

25

By using the techniques of Example 1 for the construction and fermentation, and the above isolation procedure the following subtilisin 309 variants were produced and isolated:

Position 96 insertion variants:

L96LA

L96LA + A98T + A108C + A138C

5 Position 97 insertion variants:

G97GI + S99T

Position 98 insertion variants:

A98AS + A133E + T143K

A98AT + G97D

10 A98ATGTG

A98AG

A98AS + R45K + S105G

A98AT + G97E

A98ASGTG

15 A98AP + A98G + S99A

A98AT + Y167A + R170S + A194P

A98AI+A98G+S99H+G100S+S101A

20 Position 99 insertion variants:

S99SD + S99A

S99SA

S99SE + S99T

S99SD + S99A + A133E

25 S99SD + S99A + T143K

S99SD

S99SE

S99SD + S99A + S216SP

S99SD + S99A + S216SDP

30 S99SD + S99A + P129PD

S99SD + S99A + P129PR

S99SD + S99A + L217F + A228V + A230V

S99SD + S99A + L217LP
S99SD + S99A + D42DN
S99SR + S99T
S99SQ + S99T
5 S99SD + M222S
S99SD + N76D + A194P + A230V
S99SN
S99SD + S99A + P131T

10 **EXAMPLE 3**

The "MODEL DETERGENT WASH PERFORMANCE TEST":

In order to assess the wash performance of selected subtilase variants in a standard detergent composition, standard washing experiments may be performed using the below experimental
15 conditions:

Detergent:	Model detergent
Detergent dosage	4.0 g/l
pH	10.1
20 Wash time	20 min
Temperature:	30°C
Water hardness:	15°dH
Enzyme concentration:	10 nm (in the detergent solution)
Test system:	10 ml beakers with a stirring rod
25 Textile/volume:	5 textile pieces (Ø 2.5 cm)/50 ml detergent solution
Test material:	WFK10N (egg stains)

The composition of the model detergent is as follows:

30

6.2% LAS (Nansa 80S)
2% Sodium salt of C₁₆-C₁₈ fatty acid

	4%	Non-ionic surfactant (Plurafax LF404)
	22%	Zeolite P
	10.5%	Na_2CO_3
	4%	$\text{Na}_2\text{Si}_2\text{O}_5$
5	2%	Carboxymethylcellulose (CMC)
	6.8%	Acrylate liquid CP5 40%
	20%	Sodium perborate (empirical formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$)
	0.2%	EDTA
	21%	Na_2SO_4
10		Water (balance)

pH of the detergent solution is adjusted to 10.1 by addition of HCl or NaOH. Water hardness is adjusted to 15°dH by addition of CaCl_2 and MgCl_2 ($\text{Ca}^{2+}:\text{Mg}^{2+} = 4:1$) to the test system. After
15 washing the textile pieces were flushed in tap water and air-dried.

Measurement of the reflectance (R_{variant}) on the test material is performed at 460 nm using a Macbeth ColorEye 7000 photometer
20 (Macbeth, Division of Kollmorgen Instruments Corporation, Germany). The measurements are performed accordance with the manufacturer's protocol.

In order to determine a blank value, a similar wash experiment
25 is performed without addition of enzyme. The subsequent measurement of the reflectance (R_{blank}) is performed as described right above.

A reference experiment is then performed as described above,
30 wherein the wash performance of the parent enzyme is tested. The subsequent measurement of the reflectance (R_{parent}) is performed as described right above.

The wash performance is evaluated by means of the Performance Factor (P) which is defined in accordance with the below formula:

5

$$P = (R_{\text{variant}} - R_{\text{blank}}) - (R_{\text{parent}} - R_{\text{blank}}) \\ = R_{\text{variant}} - R_{\text{parent}}.$$

Using the above test method the following results were obtained:

10

Enzyme	R (460 nm)	P
Blank (no enzyme)	40.5	-
Parent (Savinase®)	40.7	-
S99SD + S99A	43.2	2.5
S99SA	-	2.0
S99SE + S99T	-	2.0
A98AS + A133E + T143K	45.1	4.4
A98AT + G97D	-	1.5
A98ATGTG	-	1.6
A98AG	-	1.7
A98AS + R45K + S105G	-	1.8
A98AT + G97E	-	2.0
A98ASGTG	-	2.1
A98AP + A98G + S99A	-	2.3

As it appears, the subtilase variants exhibit improved wash performance on egg stains in comparison to the parent subtilase, i.e. Savinase®.

15

EXAMPLE 4

THE "OVO-INHIBITION ASSAY"

The below inhibition assay is based on the principle that the subtilase variant to be tested will catalyse the hydrolysis of a peptide-pNA bond, thereby releasing the yellow pNA, which may conveniently be followed at 405 nm. The amount of released pNA after a given period of time is a direct measure of the subtilase activity. By carrying out such hydrolysis experiments with and without inhibitor, respectively, it is possible to obtain a quantitative measure for the degree to which a certain subtilase variant is inhibited.

10

Reaction conditions:

Enzyme concentration:	0.0003 mg/ml
Conc. of trypsin inhibitor type IV-0:	0.0015 mg/ml
Initial substrate concentration:	0.81 mM
15 Reaction time:	11 min
Assay temperature:	25°C
Assay pH:	8.6
Absorbance measured at:	405 nm

20 Assay solutions:

Substrate solution (2 mM): 500 mg Suc-Ala-Ala-Pro-Phe-pNA is dissolved in 4 ml DMSO (200 mM). This solution is diluted 100 times with the buffer solution described below. The concentration of substrate in the resulting substrate solution is 2 mM.

25

Inhibitor solution (0.005 mg/ml): 5 mg trypsin inhibitor type IV-0 (Sigma T-1886) is dissolved in 10 ml water. This solution is dissolved 100 times with the buffer solution described below. The concentration of inhibitor in the resulting inhibitor solution is 0.005 mg/ml.

30

Enzyme solution (0.001 mg/ml): 1 mg enzyme is dissolved in 10 ml water. This solution is dissolved 100 times with the buffer solution described below. The concentration of enzyme in the resulting enzyme solution is 0.001 mg/ml.

5

Buffer solution (pH 8.6): 15.7 mg Tris is dissolved in an appropriate amount of water and 0.75 ml 30% (w/v) BRIJ (BRIJ 35 polyoxyethylenelaurylether, 30% (w/v), Sigma Cat. No. 430AG-6) is added. The pH is adjusted to 8.6 with 4 M NaOH and the solution is diluted to 1 liter with water.

10

Assay with inhibitor

1 volume unit (e.g. 80 μ l) inhibitor solution is mixed with 1 volume unit (e.g. 80 μ l) enzyme solution in an appropriate reaction vessel (e.g. a spectrophotometer cell or a micro titer plate) and equilibrated at 25°C for 15 min. 1.375 volume units (e.g. 110 μ l) substrate solution is added to the reaction vessel after which the absorbance at 405 nm is followed for 11 min (e.g. by measuring every 10th or 30th second). The slope of the absorbance curve is calculated using linear regression analysis. The slope of the absorbance curve is denoted $\alpha_{\text{inhibitor}}$.

20

Assay without inhibitor

1 volume unit (e.g. 80 μ l) buffer solution is mixed with 1 volume unit (e.g. 80 μ l) enzyme solution in an appropriate reaction vessel (e.g. a spectrophotometer cell or a micro titer plate) and equilibrated at 25°C for 15 min. 1.375 volume units (e.g. 110 μ l) substrate solution is added to the reaction vessel after which the absorbance at 405 nm is followed for 11 min (e.g. by measuring every 10th or 30th second). The slope of the absorbance

30

curve is calculated using linear regression analysis. The slope of the absorbance curve is denoted α .

Blank

5 1 volume unit (e.g. 80 μ l) inhibitor solution is mixed with 1 volume unit (e.g. 80 μ l) buffer solution in an appropriate reaction vessel (e.g. a spectrophotometer cell or a micro titer plate) and equilibrated at 25°C for 15 min. 1.375 volume units (e.g. 110 μ l) substrate solution is added to the reaction vessel
10 after which the absorbance at 405 nm is followed for 11 min. These measurements are not used in the calculations, but merely serve as a control that no enzyme has been added to the buffer and/or substrate solution.

15 Calculation of Residual Activity (RA)

The residual enzyme activity (RA) is calculated according to the below formula:

$$RA = (\alpha_{\text{inhibitor}}/\alpha) \times 100\%$$

20

Using the above test, the following results were obtained:

Enzyme	Residual Activity (%)
Savinase®	<5%
A98AT + Y167A + R170S + A194P	88.0
A98AI+A98G+S99H+G100S+S101A	22.0
S99SD + S99A	27.3
S99SD + S99A + A133E	39.0
S99SD + S99A + T143K	23.0
S99SD	25.0
S99SE	27.0

S99SD + S99A + S216SP	29.2
S99SD + S99A + S216SDP	35.0
S99SD + S99A + P129PD	50.0
S99SD + S99A + P129PR	21.0
S99SD+S99A+L217F+A228V+A230V	12.0
S99SD + S99A + L217LP	97.0
S99SD + S99A + D42DN	69.2
S99SR + S99T	67.7
S99SQ + S99T	25.0
S99SD + M222S	25.0
S99SD + N76D + A194P + A230V	18.4
S99SN	19.0
S99SD + S99A + P131T	35.6

As it appears, the subtilase variants were inhibited to a much smaller extent than the parent subtilase, i.e. savinase®.

5 **EXAMPLE 5**

Performance of the subtilase variant of the invention in Automatic Dishwashing (ADW)

The performance of the variant of the invention in ADW was tested in a commercial available household dishwash composition (Somat Turbo, from Henkel Washmittel GmbH) using standard conditions. The soil used was an egg/milk mixture coated on a steel plate. Further, a ballast soil containing various foodstuffs was added.

15

Detergent:	Somat Turbo
Detergent dosage	4.0 g/l
pH	10.7 (as is)
Water hardness:	3°dH (machine ion exchanger)

Temperature: 55°C
Enzyme concentration: 20 nM and 40 nM, based on the total
volume of wash water in the machine
Test method: Egg/milk soiling on steel plates as
described below
Machine: Cylinda Compact
Wash program: Program 4 without pre-flush

Materials

220 ml full cream milk
15 eggs, medium size
Steel plates, diameter 18 cm

The Somat Turbo dishwash composition was heated at 85°C for 5
minutes in a microwave oven in order to inactivate enzyme
activity in the composition.

Soiling of steel plates

220 ml full cream milk was mixed with 15 raw eggs in a Braun UK
20 kitchen machine for 2 minutes, After sieving, stainless steel
plates were soiled in the mixture by immersion.

The plates were dried overnight at room temperature in an
upright position. The dried plates were then heated at 120°C for
25 45 minutes in order to denature the proteins on the surface.

ADW experiments

For each experiment, 10 soiled plates were washed without pre-
wash (Program 4) in a Cylinda Compact machine. In addition to
30 the soiled plates, the machine was filled up with 10 porcelain
plates, 4 glasses, 4 cups and 16 pieces of cutlery.

Furthermore, 50 g of ballast slurry was added to the machine.
The composition of the slurry was as follows:

Potato starch (5.43%), wheat flour (4.38%), vegetable oil
5 (4.32%), margarine (4.32%), lard (4.32%), cream (8.76%), full
cream milk (8.76%), eggs (8.76%), tomato ketchup (3.00%),
barbecue sauce (2.19%), mustard (4.00%), benzoic acid (0.73%),
water (3 mM Ca^{2+} + Mg^{2+}) (36.71%).

10 Measurements and calculations

The light reflection values (R-values) were measured at six
different locations on the plates using a Minolta Chroma Meter
(Type: CR-300). Measurements were made on clean plates (R_{clean}),
on soiled plates after heating (R_{soiled}) and on plates after wash
15 ($R_{\text{after wash}}$).

The removed protein film (%RPF) was calculated according to the
below formula:

$$20 \quad \%RPF = 100\% \times (R_{\text{after wash}} - R_{\text{soiled}}) / (R_{\text{clean}} - R_{\text{soiled}})$$

Using the above test method the following results were obtained
(\pm indicates the standard deviation):

Enzyme	%RPF (20 nM)	%RPF (40 nM)
Savinase®	3.9 \pm 1.6	3.0 \pm 1.0
S99SD + S99A	13.8 \pm 5.2	77.1 \pm 2.2

25

As it appears, the variant of the invention has a superior
performance as compared to Savinase®.

EXAMPLE 6

Wash performance of the subtilase variant of the invention in a commercially available powder detergent

- 5 In order to assess the wash performance of selected subtilase variants in a commercial detergent composition, standard washing experiments were performed using the below experimental conditions:

90

Detergent dosage: 4 g/l
Wash temperature: 30°C
Washing time: 20 minutes
Water hardness: 15°dH ($\text{Ca}^{2+}:\text{Mg}^{2+} = 4:1$)
5 pH: Not adjusted
Enzyme concentrations: 1, 2, 5, 10, 30 nM
Test system: 150 ml glass beakers with a
stirring rod
Textile/volume: 5 textile pieces (Ø 2.5 cm) in 50
10 ml detergent
Test material: WFK10N (egg stains)

The detergent used was obtained from supermarket in Germany
(Persil Megapearls). Prior to use all enzymatic activity was in
15 the detergents were inactivated by microwave treatment (5
minutes, 85°C).

The reflectance measurements were performed as described in
Example 3 herein.

20

The data (the R values) were evaluated as follows:

A variant having a higher R-value than savinase® was given the
value 1.

25

A variant having a lower R-value than savinase® was given the
value -1.

A variant having a R-value similar to savinase® was given the
30 value 0.

Results:

	<u>Variant</u>	<u>Value</u>
	Savinase®	0
5	L96LA	1
	L96LA+A98T+A108C+A138C	1
	<u>G97GI+S99T</u>	<u>1</u>

As I appears, the subtilase variants exhibit improved wash
10 performance in a commercial detergent as compared to savinase®.

CLAIMS

1. Use of a subtilase variant for removal of egg stains from laundry or from hard surfaces, the subtilase variant comprising at least one additional amino acid residue in the active site loop (b) region from position 95 to 103 (BASBPN numbering).
2. Use according to claim 1, wherein the additional amino acid residue has been inserted in a position selected from the group consisting of: between positions 95 and 96, between positions 96 and 97, between positions 97 and 98, between positions 98 and 99, between positions 99 and 100, between positions 100 and 101, between positions 101 and 102, between positions 102 and 103, between position 103 and 104 and combinations thereof.
3. Use according to claim 2, wherein the additional amino acid residue has been inserted in a position selected from the group consisting of: between positions 98 and 99 and between positions 99 and 100.
4. Use according to any of claims 1-3, where the variant - when tested in the "Ovo-inhibition Assay" disclosed in Example 4 herein - has a Residual Activity of at least 10%, such as at least 15%, preferably at least 20%, more preferably at least 25%.
5. Use according to any of claims 1-4, wherein the insertion between positions 98 and 99 is selected from the group consisting of X98XA, X98XT, X98XG and X98XS.
6. Use according to any of claims 1-4, wherein the insertion between positions 99 and 100 is selected from the group consisting of X99XD, X99XE, X99XK and X99XR.

7. Use according to any of the preceding claims, wherein the variant comprises at least one further modification.

8. Use according to claim 7, wherein the further modification is performed
5 in a position selected from the group consisting of: substitution in position 99, substitution in position 133, substitution in position 143, substitution in position 167, substitution in position 170, substitution in position 194, insertion between positions 42 and 43, insertion between positions 129 and 130, insertion between positions 216 and 217, insertion
10 between 217 and 218, and combinations thereof.

9. Use according to claim 8, wherein the variant is selected from the group consisting of

15 a variant comprising an insertion of at least one additional amino acid residue between positions 98 and 99 and further comprising a substitution in positions 133 and 143,

a variant comprising an insertion of at least one additional amino acid
20 residue between positions 99 and 100 and further comprising a substitution in position 99,

a variant comprising an insertion of at least one additional amino acid residue between positions 98 and 99 and further comprising substitutions
25 in positions 167, 170 and 194,

a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising an insertion of at least one additional amino acid residue between positions 216 and
30 217,

a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising an insertion of at least one additional amino acid residue between positions 217 and 218,

5

a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising an insertion of at least one additional amino acid residue between positions 42 and 43, and

10

a variant comprising an insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising an insertion of at least one additional amino acid residue between positions 129 and 130.

15

10. Use according to any of the preceding claims, wherein the parent subtilase belongs to the sub-group I-S1.

11. Use according to claim 10, wherein the parent subtilase is
20 selected from the group consisting of BSS168, BASBPN, BSSDY, and BLSCAR, or functional variants thereof having retained the characteristic of sub-group I-S1.

12. Use according to any of claims 1-9, wherein the parent
25 subtilase belongs to the sub-group I-S2.

13. Use according to claim 12, wherein the parent subtilase is
selected from the group consisting of BLS147, BLSAVI, BAPB92,
TVTHER and BYSYAB, or functional variants thereof having
30 retained the characteristic of sub-group I-S2.

14. Use according to claim 13, wherein the parent subtilase is
BLSAVI (SEQ ID NO: 1).

15. Use according to claim 14, wherein the variant is S99SD +
5 S99A.

16. Use according to claim 14, wherein the variant is S99SR +
S99T.

10 17. Use according to claim 14, wherein the variant is A98AS +
A133E + T143K.

18. Use according to claim 14, wherein the variant is A98AT +
Y167A + R170S + A194P.

15 19. Use according to claim 14, wherein the variant is S99SD +
S99A + P129PD.

20 20. Use according to claim 14, wherein the variant is S99SD +
S99A + S216SP.

21. Use according to claim 14, wherein the variant is S99SD +
S99A + S216SDP.

25 22. Use according to claim 14, wherein the variant is S99SD +
S99SA + L217LP.

23. Use according to claim 14, wherein the variant is S99SD +
D42DN.

30 24. Use according to claim 14, wherein the variant is S99SD +
S99A + D42DN.

25. A subtilase variant selected from the group consisting of

5 a variant comprising at least one additional amino acid residue in the active site (b) loop corresponding to the insertion of at least one additional amino acid residue between positions 98 and 99 and further comprising at least one additional modification (BASBPN numbering), and

10 a variant comprising at least one additional amino acid residue in the active site (b) loop corresponding to the insertion of at least one additional amino acid residue between positions 99 and 100 and further comprising at least one additional modification (BASBPN numbering),

15 where the variant - when tested in the "Ovo-inhibition Assay" disclosed in Example 4 herein - has a Residual Activity of at least 10%.

26. The variant according to claim 25, where the variant has a Residual Activity of at least 15%, preferably at least 20%, more preferably at least 25%.

27. The variant according to claims 25 and 26 having the characteristic as defined in any of claims 9-24

25 28. A subtilase variant as defined in any of claims 9-24.

29. An isolated DNA sequence encoding a subtilase variant as defined in any of claims 25-28.

30 30. An expression vector comprising the isolated DNA sequence of claim 29.

31. A microbial host cell transformed with the expression vector of claim 30.

32. A microbial host cell according to claim 31, which is a bacterium,
5 preferably a *Bacillus*, especially a *B. lentus*.

33. A microbial host cell according to claim 31, which is a fungus or yeast, preferably a filamentous fungus, especially an *Aspergillus*.

10 34. A method for producing a subtilase variant according to any of claims 25-27, wherein a host according to any claims 31-33 is cultured under conditions conducive to the expression and secretion of said variant, and the variant is recovered.

15 35. A cleaning or detergent composition, preferably a laundry or dishwash composition, comprising the variant according to any of claims 25-28.

36. A composition according to claim 35, which additionally
20 comprises a cellulase, a lipase, a cutinase, an oxidoreductase, another protease, an amylase or a mixture thereof.

37. Use of a variant as defined in any of claims 25-28 in a cleaning or detergent composition, preferably a laundry and/or a
25 dishwash composition.

38. A method for removal of egg stains from a hard surface or from laundry, the method comprising contacting the egg stain-containing hard surface or the egg stain-containing laundry with
30 a cleaning or detergent composition, preferably a laundry or dishwash composition, containing a subtilase variant comprising

at least one additional amino acid residue in the active site loop (b) region from position 95 to 103 (BASBPN numbering).

39. A method according to claim 38, wherein the variant has the
5 characteristics as defined in any of claims 2-24.

40. A method according to any of claims 38-39, wherein the composition additionally comprises a cellulase, a lipase, a cutinase, an oxidoreductase, another protease, an amylase or a
10 mixture thereof.

41. Use of a cleaning or detergent composition, preferably a laundry or dishwash composition, containing a subtilase variant comprising at least one additional amino acid residue in the active site loop
15 (b) region from position 95 to 103 (BASBPN numbering) for removal of egg stains from laundry or from hard surfaces.

42. Use according to claim 41, wherein the variant has the characteristics as defined in any of claims 2-24.

20

43. Use according to any of claims 41-42, wherein the composition additionally comprises a cellulase, a lipase, a cutinase, an oxidoreductase, another protease, an amylase or a mixture thereof.

25

1/1

No: 1 10 20 30 40 50
a) AQSVPYGVSQIKAPALHSQGYTGSNVKVAVIDSGIDSSHPLDKVAGGASM
5 b) AQSVPWGISRVQAPAAHNRGLTGSGVKVAVLDTGI*STHPDLNIRGGASF

No: 60 70 80 90 100
a) VPSETNPFQDNNSHGTHVAGTVAALNNSIGVLGVAPSASLYAVKVLGADG
b) VPGEPT*QDGNHGHGTHVAGTIAALNNSIGVLGVAPSAELYAVKVLGASG
10

No: 110 120 130 140 150
a) SGQYSWIINGIEWAIANNMDVINMSLGGPSGSAALKAAVDKAVASGVVVV
b) SGSVSSIAQGLEWAGNNGMHVANLSLGSPSPSATLEQAVNSATSRGVLVV

15 No: 160 170 180 190 200
a) AAAGNEGTSGSSSTVGYPGKYPSVIAVGAVDSSNQRASFSSVGPELDVMA
b) AASGNSG*AGS***ISYPARYANAMAVGATDQNNNRASFSSQYGAGLDIVA

No: 210 220 230 240 250
20 a) PGVSIQSTLPGNKYGAYNGTSMASPHVAGAAALILSKHPNWTNTQVRSSL
b) PGVNVQSTYPGSTYASLNGTSMATPHVAGAAALVKQKNPSWSNVQIRNHL

No: 260 270 275
a) ENTTTKLGDSFYYGKGLINVQAAAQ
25 b) KNTATSLGSTNLYGSGLVNAEAAATR

Fig.1

SEQUENCE LISTING

<110> Novozymes A/S

<120> Subtilase variants having an improved performance on
egg stains

<130> 6108.204-WO

<140>

<141>

<160> 105

<170> PatentIn Ver. 2.1

<210> 1

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Nomenclature
example

<400> 1

Gly Gly Lys Ala Ser
1 5

<210> 2

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 2

cagaagatgt ggacgcgctt g

21

<210> 3

<211> 50

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 3

ctgcacgttt accccgggtg cgacaatgtc aaggcctggg ccatactgtg 50

<210> 4

<211> 52

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 4

ctcgatacag ggatatccac tcatccagat ctaaacaata ttcgtggtgg cg 52

<210> 5

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 5

ccgaacctga accatccgcg gcccctagga ctttaacagc 40

<210> 6

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 6

aaccgcacag cgttttttta ttgattaacg cgttgc 36

<210> 7

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 7

tgaaccgctg gtggggccta ggactttaac ag

32

<210> 8

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 8

gattaacgcg ttgccgcttc tg

22

<210> 9

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 9

cagaagatgt ggacgcgctt g

21

<210> 10

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 10

gaccgaacct gaacctgag tggcgcttag gac

33

<210> 11
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 11
gattaacgcg ttgccgcttc tg 22

<210> 12
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sense primer

<400> 12
gagttaagcc cagaagatgt ggacgcg 27

<210> 13
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 13
gaccgaacct gaaccatcgc tcgcccctag gac 33

<210> 14
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 14

aggagtagcc gacgatgtac cgtttaa

27

<210> 15

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 15

gagttaagcc cagaagatgt ggacgcg

27

<210> 16

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Nomenclature
example

<400> 16

Ala Gly Lys Ala Ser Leu

1

5

<210> 17

<211> 70

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 17

ccattccaat ccctggcaaa tcgagctgac cgaacctgaa ccgctggtag ccgctaggac 60
tttaacagcg 70

<210> 18

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 18

aacgcctcta gaagtcgcgc tattaacaca ttgctcgagt gtgg

44

<210> 19

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 19

aaccgcacag cgttttttta ttgattaacg cgttgc

36

<210> 20

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 20

cagaagatgt ggacgcgctt g

21

<210> 21

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 21

aaccgctggt ggcgtctagg actttaacag cg

32

<210> 22
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 22
gattaacgcg ttgccgcttc tg 22

<210> 23
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sense primer

<400> 23
cagaagatgt ggacgcgctt g 21

<210> 24
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 24
aaccgctggt ggcttctagg actttaacag cg 32

<210> 25
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 25

gattaacgcg ttgccgttc tg

22

<210> 26

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 26

gagttaagcc cagaagatgt ggacgcg

27

<210> 27

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Nomenclature
example

<400> 27

Ala Gly Gly Leu

1

<210> 28

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 28

accgaacctg aacctgcgct cgcccctagg

30

<210> 29

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 29

cagaagatgt ggacgcgctt g

21

<210> 30

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 30

gaccgaacct gagccctcgg tggcgcctag gac

33

<210> 31

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 31

gattaacgcg ttgccgcttc tg

22

<210> 32

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 32

cccttcgcca agtgagactc tcgagcaagc tg

32

<210> 33

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 33

acagcgtttt tttattgatt aacgcgttgc

30

<210> 34

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 34

aaagtcttag gggccgccga cggttcaggt tcggtcagc

39

<210> 35

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 35

gagttaagcc cagaagatgt ggacgcg

27

<210> 36

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 36

tggttaatagc gcgaaatcca gaggcgttct tg

32

<210> 37

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 37

acagcgtttt tttattgatt aacgcgttgc

30

<210> 38

<211> 275

<212> PRT

<213> B. amyloliquefaciens (subtilisin BPN')

<400> 38

Ala Gln Ser Val Pro Tyr Gly Val Ser Gln Ile Lys Ala Pro Ala Leu

1

5

10

15

His Ser Gln Gly Tyr Thr Gly Ser Asn Val Lys Val Ala Val Ile Asp

20

25

30

Ser Gly Ile Asp Ser Ser His Pro Asp Leu Lys Val Ala Gly Gly Ala

35

40

45

Ser Met Val Pro Ser Glu Thr Asn Pro Phe Gln Asp Asn Asn Ser His

50

55

60

Gly Thr His Val Ala Gly Thr Val Ala Ala Leu Asn Asn Ser Ile Gly

65

70

75

80

Val Leu Gly Val Ala Pro Ser Ala Ser Leu Tyr Ala Val Lys Val Leu

85

90

95

Gly Ala Asp Gly Ser Gly Gln Tyr Ser Trp Ile Ile Asn Gly Ile Glu

100

105

110

Trp Ala Ile Ala Asn Asn Met Asp Val Ile Asn Met Ser Leu Gly Gly

115

120

125

Pro Ser Gly Ser Ala Ala Leu Lys Ala Ala Val Asp Lys Ala Val Ala

130

135

140

Ser Gly Val Val Val Val Ala Ala Ala Gly Asn Glu Gly Thr Ser Gly

145

150

155

160

Ser Ser Ser Thr Val Gly Tyr Pro Gly Lys Tyr Pro Ser Val Ile Ala

	165	170	175
Val Gly Ala Val Asp Ser Ser Asn Gln Arg Ala Ser Phe Ser Ser Val			
	180	185	190
Gly Pro Glu Leu Asp Val Met Ala Pro Gly Val Ser Ile Gln Ser Thr			
	195	200	205
Leu Pro Gly Asn Lys Tyr Gly Ala Tyr Asn Gly Thr Ser Met Ala Ser			
	210	215	220
Pro His Val Ala Gly Ala Ala Ala Leu Ile Leu Ser Lys His Pro Asn			
	225	230	235 240
Trp Thr Asn Thr Gln Val Arg Ser Ser Leu Glu Asn Thr Thr Thr Lys			
	245	250	255
Leu Gly Asp Ser Phe Tyr Tyr Gly Lys Gly Leu Ile Asn Val Gln Ala			
	260	265	270
Ala Ala Gln			
	275		

<210> 39

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 39

aaagtcctag gggccgccga cggttcaggt tcggtcagc

39

<210> 40

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 40

gagttaagcc cagaagatgt ggacgcg

27

<210> 41

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 41

gagttaagcc cagaagatgt ggacgcg

27

<210> 42

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 42

ccgaacctga accatccgcg gcccttagga cttaacagc

40

<210> 43

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 43

gatgtaccgt ttaaagggt ggcataatgtt gaacc

35

<210> 44

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 44
aaccgcacag cgttttttta ttgattaacg cgttgc 36

<210> 45
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sense primer

<400> 45
gagttaagcc cagaagatgt ggacgcg 27

<210> 46
<211> 40
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 46
ccgaacctga accatccgcg gcccttagga ctttaacagc 40

<210> 47
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 47
gtaccgttta aaggatcgct ggcataatggt gaacc 35

<210> 48
<211> 36
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 48

aaccgcacag cgttttttta ttgattaacg cgttgc

36

<210> 49

<211> 269

<212> PRT

<213> Bacillus lentus (subtilisin 309)

<400> 49

Ala Gln Ser Val Pro Trp Gly Ile Ser Arg Val Gln Ala Pro Ala Ala
1 5 10 15

His Asn Arg Gly Leu Thr Gly Ser Gly Val Lys Val Ala Val Leu Asp
20 25 30

Thr Gly Ile Ser Thr His Pro Asp Leu Asn Ile Arg Gly Gly Ala Ser
35 40 45

Phe Val Pro Gly Glu Pro Ser Thr Gln Asp Gly Asn Gly His Gly Thr
50 55 60

His Val Ala Gly Thr Ile Ala Ala Leu Asn Asn Ser Ile Gly Val Leu
65 70 75 80

Gly Val Ala Pro Ser Ala Glu Leu Tyr Ala Val Lys Val Leu Gly Ala
85 90 95

Ser Gly Ser Gly Ser Val Ser Ser Ile Ala Gln Gly Leu Glu Trp Ala
100 105 110

Gly Asn Asn Gly Met His Val Ala Asn Leu Ser Leu Gly Ser Pro Ser
115 120 125

Pro Ser Ala Thr Leu Glu Gln Ala Val Asn Ser Ala Thr Ser Arg Gly
130 135 140

Val Leu Val Val Ala Ala Ser Gly Asn Ser Gly Ala Gly Ser Ile Ser
145 150 155 160

Tyr Pro Ala Arg Tyr Ala Asn Ala Met Ala Val Gly Ala Thr Asp Gln
165 170 175

Asn Asn Asn Arg Ala Ser Phe Ser Gln Tyr Gly Ala Gly Leu Asp Ile
180 185 190

Val Ala Pro Gly Val Asn Val Gln Ser Thr Tyr Pro Gly Ser Thr Tyr
 195 200 205

Ala Ser Leu Asn Gly Thr Ser Met Ala Thr Pro His Val Ala Gly Ala
 210 215 220

Ala Ala Leu Val Lys Gln Lys Asn Pro Ser Trp Ser Asn Val Gln Ile
 225 230 235 240

Arg Asn His Leu Lys Asn Thr Ala Thr Ser Leu Gly Ser Thr Asn Leu
 245 250 255

Tyr Gly Ser Gly Leu Val Asn Ala Glu Ala Ala Thr Arg
 260 265

<210> 50

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 50

gagttaagcc cagaagatgt ggacgcg

27

<210> 51

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
 primer

<400> 51

ccgaacctga accatccgcg gccctagga ctttaacagc

40

<210> 52

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 52

gtgtggcact tggcgagtca gggcttccta aactc

35

<210> 53

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 53

aaccgcacag cgttttttta ttgattaacg cgttgc

36

<210> 54

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 54

gagttaagcc cagaagatgt ggacgcg

27

<210> 55

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 55

ccgaacctga accatccgcg gcccttagga cttaacagc

40

<210> 56

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 56

gtgtggcact tggcgatcga gggcttccta aactc

35

<210> 57

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 57

aaccgcacag cgttttttta ttgattaacg cgttgc

36

<210> 58

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 58

gagttaagcc cagaagatgt ggacgcg

27

<210> 59

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 59

ccgaacctga accatccgcg gccccctagga ctttaacagc

40

<210> 60

<211> 13222

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: pJS3: E. coli

- B. subtilis shuttle vector

<400> 60

```

aattccggcc caacgatggc tgatttccgg gttgacggcc ggcggaacca aggggtgata 60
ggtcggcgga aatgaaggcc tgcggcgagt gcgggccttc ttaaggccgg gttgctaccg 120
actaaaggcc caactgccgg cgccttggt tccccactag ccagccgcct ttacttccgg 180
acgccgctca cgcccgaag tgttttgagg attataatca gagtatattg aaagtttcgc 240
gatcttttgc tataattggt ttaggcatag tgcaatcgat tgtttgagaa aagaagaaga 300
acaaaactcc taatattagt ctcatataac tttcaaagcg ctagaaaagc atattaacaa 360
aatccgtatc acgtagctta acaaactctt ttcttcttct ccataaaaat acctgtctcg 420
tcatcagaca gggatatttt tatgctgtcc agactgtccg ctgtgtaaaa ataaggaata 480
aagggggggt gttattattt ggtattttta tggaacagac agtagtctgt cccataaaaa 540
atacgacagg tctgacaggc gacacatttt tattccttat tccccccaa caataataaa 600
tactgatatg taaaatataa tttgtataag aaaatgagag ggagaggaaa catgattcaa 660
aaacgaaagc ggacagtttc gttcagactt gtgcttatgt atgactatac attttatatt 720
aaacatatct ttttactctc cctctccttt gtactaagtt tttgctttcg cctgtcaaag 780
caagtctgaa cacgaatata gcacgtggt atttgtcagt ttgccgatta caaaaacatc 840
agccgtaaat ggcacgctga tgcagtattt tgaatggtat acgccgaacg acggccagca 900
cgtgcgacaa taaacagtca aacggctaatt gttttttagt tcggcattta ccgtgcgact 960
acgtcataaa acttaccata tgcggcttgc tgcgggtcgt ttggaaacga ttgcagaatg 1020
atgcggaaca tttatcggat taacttaacg ttaatatattg tttcccaata ggcaaatctt 1080
tctaactttg atacgtttta aacctttgct aacgtcttac tacgccttgt aaatagccta 1140
attgaattgc aattataaac aaagggttat ccgtttagaa agattgaaac tatgcaaat 1200
actaccagct tggacaagtt ggtataaaaa tgaggaggga aaccgaatga agaaaccgtt 1260
ggggaaaatt gtcgaagca ccgcactact catttctggt tgatggtcga acctgttcaa 1320
ccatattttt actcctccct ttggcttact tctttggcaa ccccttttaa cagcgttcgt 1380
ggcgtgatga gtaaagacaa gcttttagtt catcgatcgc atcggctgct gaagaagcaa 1440
aagaaaaata tttaattggc tttaatgagc aggaagctgt cagtgaagtt gtagaacaag 1500
cgaaaatcaa gtagctagcg tagccgacga cttcttcggt ttctttttat aaattaaccg 1560
aaattactcg tccttcgaca gtcactcaaa catcttggtc tagaggcaaa tgacgaggtc 1620
gccattctct ctgaggaaga ggaagtcgaa attgaattgc ttcataaatt tgaaacgatt 1680
cctgttttat ccgttgaggt atctccggtt actgctccag cggtaagaga gactccttct 1740
ccttcagctt taacttaacg aagtacttaa actttgctaa ggacaaaata ggcaactcaa 1800
aagcccagaa gatgtggacg cgcttgaact cgatccagcg atttcttata ttgaagagga 1860
tgcagaagta acgacaatgg cgcaatcggg accatgggga ttcgggtctt ctacacctgc 1920
gcgaacttga gtaggtcgc taaagaatat aacttctcct acgtcttcat tgctgttacc 1980
gcgttagcca tggtagccct attagccgtg tgcaagcccc agctgcccac aaccgtggat 2040
tgacaggttc tgggtgaaaa gttgctgtcc tcgatacagg gatattccact catccagatc 2100
taatcgccac acgttcgggg tgcacgggta ttggcaccta actgtccaag accacatttt 2160
caacgacagg agctatgtcc ctataggtga gtaggtctag taaatatctg tgggtggcgca 2220

```

```

agctttgtac caggggaacc gtcgactcaa gatgggaatg ggcattggcac gcatgtggcc 2280
gggacgatcg ctgctttaaa atttataagc accaccgctg tcgaaacatg gtcccccttg 2340
cagctgagtt ctacccttac cgtaccgtg cgtacaccgg ccctgctagc gacgaaat 2400
caattcgatt ggcggtcttg gcgtagctcc tagcgctgag ctatacgctg ttaaagtcct 2460
aggggcgagc gggtcagggt cggtcagctc gattgcccga gtttagctaa ccgcaagaac 2520
cgcacgaggg atcgcgactc gatatgcgac aatttcagga tccccgctcg ccaagtccaa 2580
gccagtcgag ctaacggggt ggattggaat gggcagggaa caatggcatg cacgttgcta 2640
at ttgagttt aggaagccct tcgccaagtg ccacactcga gcaagctgtt aatagcgcg 2700
cctaaccctta cccgtccctt gttaccgtac gtgcaacgat taaactcaaa tccttcggga 2760
agcggttcac gggtgtgagct cgttcgacaa ttatcgctc cttctagagg cgttctgtt 2820
gtagcggcac ctgggaattc aggtgcaggc tcaatcagct atccggcgcg ctatgcgaac 2880
gcaatggcag tcggagctac gaagatctcc gcaagaacaa catcgccgta gacccttaag 2940
tccacgtccg agttagtcga taggcgcgcg gatacgcttg cgttaccgtc agcctcgatg 3000
tgatcaaaac aacaaccgcg ctagcttttc acagtatggc gcaggccttg acattgtcgc 3060
accgggggta aacgtgcaga gcacataccc aggttcaaca actagttttg ttgttggcgc 3120
gatcgaaaag tgcataccg cgtccggaac tgtaacagcg tgggccccat ttgcacgtct 3180
cgtgtatggg tccaagttgt tatgccagct taaacggtag atcgatggct actcctcatg 3240
ttgcaggtgc ggccgccctt gttaaacaaa agaaccatc ttggtctaata gtacaaattc 3300
atacggtcga at ttgccatg tagctaccga tgaggagtag aacgtccacg ccggcgggaa 3360
caatttgttt tcttgggtag aaccagatta catgtttaag gaaatcatct aaagaatacg 3420
gcaactagtt taggaagcac gaacttgat ggaagcggac ttgttaacgc agaagcggca 3480
acgcgttaat caataaaaaa ctttagtaga tttcttatgc cgttgatcaa atccttcgtg 3540
cttgaacata ccttcgcctg aacaattgcg tcttcgccgt tgcgcaatta gttatTTTTT 3600
acgtgtgtcg gttaaagggc acagcgtttt tttgtgtatg gatcagcttg gcgtaatcat 3660
ggtcatagct gtttcctgtg tgaaattgtt atccgctcac tgcgacacgc caatttcccg 3720
tgtcgaaaaa aaacacatac ctagtgaac cgcattagta ccagtatcga caaaggacac 3780
actttaacaa taggcgagtg aattccacac aacatacgag ccggaagcat aaagtgtaaa 3840
gcctgggggtg cctaatagag gagctaactc acattaattg cgttgcgctc actgcccgtc 3900
ttaagggtgtg ttgtatgctc ggccttcgta tttcacattt cggacccac ggattactca 3960
ctcgattgag tgtaattaac gcaacgcgag tgacgggcga ttccagtcgg gaaacctgtc 4020
gtgccagctg cattaatgaa tcggccaacg cgcggggaga ggcgggttgc gtattgggcg 4080
ctcttcgcgt tcctcgctca aaggtcagcc ctttgacag cacggctgac gtaattactt 4140
agccggttgc gcgccctct ccgcaaacg cataaccgc gagaaggcga aggagcgagt 4200
ctgactcgct gcgctcggtc gttcggctgc ggcgagcgg atcagctcac tcaaaggcgg 4260
taatacgggt atccacagaa tcaggggata acgcaggaaa gactgagcga cgcgagccag 4320
caagccgacg ccgctcgcca tagtcgagtg agtttccgcc attatgcaa taggtgtctt 4380
agtccccat tgctgccttt gaacatgtga gcaaaaggcc agcaaaaggc caggaaccgt 4440
aaaaaggcgg cgttgctggc gtttttccat aggtccgcc cccctgacga gcatcacaaa 4500
cttgtaact cgttttccgg tcgtttccg gtccttgga tttttccggc gcaacgaccg 4560
caaaaaggta tccgaggcgg ggggactgct cgtagtgtt aatcgacgt caagtcagag 4620
gtggcgaaac ccgacaggac tataaagata ccaggcgtt cccctggaa gctccctcgt 4680
gcgctctcct gttccgacct ttagctgcga gttcagctc caccgctttg ggctgtcctg 4740
atatttctat ggtccgcaa gggggacct cgaggagca cgcgagagga caaggctggg 4800
tgccgcttac cggatacctg tccgccttt ccccttcggg aagcgtggcg ctttctcata 4860
gtcacgctg taggtatctc agttcggtgt aggtcgttcg acggcgaatg gcctatggac 4920
aggcggaaag agggaagccc ttcgcaccgc gaaagagtat cgagtgcgac atccatagag 4980
tcaagccaca tccagcaagc ctccaagctg ggctgtgtgc acgaaccccc cgttcagccc 5040
gaccgctgcg ccttatccgg taactatcgt cttgagttca acccggttaag acacgactta 5100

```



```

gaggttcgac ccgacacacg tgcttggggg gcaagtcggg ctggcgacgc ggaataggcc 5160
attgatagca gaactcaggt tgggccattc tgtgctgaat tcgccactgg cagcagccac 5220
tggtaacagg attagcagag cgaggtatgt aggcggtgct acagagttct tgaagtgggtg 5280
gcctaactac ggctacacta agcggtgacc gtcgtcgggt accattgtcc taatcgtctc 5340
gtcccataca tccgccacga tgtctcaaga acttcaccac cggattgatg ccgatgtgat 5400
gaaggacagt atttggtatc tgcgctctgc tgaagccagt taccttcgga aaaagagttg 5460
gtagctcttg atccggcaaa caaaccaccg ctggtagcgg ctccctgtca taaaccatag 5520
acgcgagacg acttcggtca atggaagcct tttctcaac catcgagaac taggccgttt 5580
gtttgggtggc gaccatcgcc tggttttttt gtttgcaagc agcagattac gcgcagaaaa 5640
aaaggatctc aagaagatcc tttgatcttt tctacggggt ctgacgctca gtggaacgaa 5700
acaaaaaaa caaacgttcg tcgtctaatt cgcgctcttt tttcctagag ttcttctagg 5760
aaactagaaa agatgcccc aactgcgagt caccttgctt aactcacgtt aagggaattt 5820
ggcatagaga ttatcaaaaa ggatcttcac ctatgcctt ttaaattaaa aatgaagttt 5880
taaatacaatc taaagtatat ttgagtgcga ttcctaaaa ccagtactct aatagttttt 5940
cctagaagtg gatctaggaa aatttaattt ttacttcaaa atttagttag atttcatata 6000
atgagtaaac ttggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga 6060
tctgtctatt tcgttcaccc atagttgcct gactccccgt tactcatttg aaccagactg 6120
tcaatggtta cgaattagtc actccgtgga tagagtcgct agacagataa agcaagtagg 6180
tatcaacgga ctgaggggca cgtgtagata actacgatac gggagggcct accatctggc 6240
cccagtgtg caatgatacc gcgagacca cgctcaccgg ctccagattt atcagcaata 6300
gcacatctat tgatgctatg ccctcccgaa tggtagaccg gggtcacgac gttactatgg 6360
cgctctgggt gcgagtggcc gaggtctaaa tagtcgttat aaccagccag ccggaagggc 6420
cgagcgcaga agtggtcctg caactttatc cgctccatc cagtctatta attgttgccg 6480
ggaagctaga gtaagtagtt ttggtcggtc ggccttccc gctcgcgtct tcaccaggac 6540
gttgaaatag gcggaggtag gtcagataat taacaacggc ctttcgatct cattcatcaa 6600
cgccagttaa tagtttgcc aacgttggtg ccattgctac aggcacgtg gtgtcacgct 6660
cgctgtttg tatggcttca ttcagctccg gttcccaac gcggtcaatt atcaaacgcg 6720
ttgcaacaac ggtaacgatg tccgtagcac cacagtgcga gcagcaaacc ataccgaagt 6780
aagtcgaggc caagggttgc atcaaggcga gttacatgat ccccatgtt gtgcaaaaaa 6840
gcggttagct ccttcggtcc tccgatcgtt gtcagaagta agttggccgc agtggttatca 6900
tagttccgct caatgtacta gggggtacaa cacgtttttt cgccaatcga ggaagccagg 6960
aggctagcaa cagtcttcat tcaaccggcg tcacaatagt ctcatggtta tggcagcact 7020
gcataattct ctactgtca tgccatccgt aagatgcttt tctgtgactg gtgagtactc 7080
aaccaagtca ttctgagaat gagtaccaat accgtcgtga cgtattaaga gaatgacagt 7140
acggtaggca ttctacgaaa agacactgac cactcatgag ttggttcagt aagactctta 7200
agtgtatgcg gcgaccgagt tgctcttgcc cggcgtcaat acgggataat accgcgccac 7260
atagcagaac tttaaaagtg ctcatcattg gaaaacgttc tcacatacgc cgctggctca 7320
acgagaacgg gccgcagtta tgccctatta tggcgcgggt tatcgtcttg aaattttcac 7380
gagtagtaac cttttgcaag ttcggggcga aaactctcaa ggatcttacc gctgttgaga 7440
tccagttcga tgtaaccac tcgtgcaccc aactgatctt cagcatcttt tactttcacc 7500
aagccccgct tttgagagtt cctagaatgg cgacaactct aggtcaagct acattgggtg 7560
agcacgtggg ttgactagaa gtcgtagaaa atgaaagtgg agcgtttctg ggtgagcaaa 7620
aacaggaagg caaaatgccg caaaaaaggg aataaggcg acacggaaat gttgaatact 7680
catactcttc ctttttcaat tcgcaaagac ccactcgttt ttgtccttcc gttttacggc 7740
gttttttccc ttattcccgc tgtgccttta caacttatga gtatgagaag gaaaaagtta 7800
gatcctctac gccggacgca tcgtggccgg catcaccggc gccacagggt cggttgctgg 7860
cgcctatata gccgacatca ccgatgggga agatcgggct ctaggagatg cggcctgcgt 7920
agcaccggcc gtagtggccg cgggtgtccac gccaacgacc gcgatatag cggctgtagt 7980

```

```

ggctaccctt tctagccga cgccacttcg ggctcatgag cgcttgtttc ggctgggta 8040
tgggtggcagg cccgtggccg ggggactggt gggcgccatc tccttgcatg ccttttagtc 8100
gcggtgaagc ccgagtaact gcgaacaaag ccgcacccat accaccgtcc gggcaccggc 8160
cccctgacaa ccgcggttag aggaacgtac ggaaaatcag cagctgattt cactttttgc 8220
attctacaaa ctgcataact catatgtaaa tcgctccttt ttaggtggca caaatgtgag 8280
gcattttcgc tctttccggc gtcgactaaa gtgaaaaacg taagatggtt gacgtattga 8340
gtatacattt agcgaggaaa aatccaccgt gtttactctc cgtaaaagcg agaaaggccg 8400
gaggctagtt acccttaagt tattggtatg actggtttta agcgcaaaaa aagttgcttt 8460
ttcgtaccta ttaatgtatc gttagaaaac cgactgtaaa ctccgatcaa tgggaattca 8520
ataaccatac tgacaaaaat tcgctgtttt ttcaacgaaa aagcatggat aattacatag 8580
caatcttttg gctgacattt aagtacagtc ggcattatct catattataa aagccagtca 8640
ttaggcctat ctgacaattc ctgaatagag ttcataaaca atcctgcatg ataaccatca 8700
ttcatgtcag ccgtaataga gtataatatt ttcggtcagt aatccggata gactgttaag 8760
gacttatctc aagtatttgt taggacgtac tattggtagt caaacagaat gatgtacctg 8820
taaagatagc ggtaaatata ttgaattacc tttattaatg aattttcctg ctgtaataat 8880
gggtagaagg taattactat gtttgtctta ctacatggac atttctatcg ccattttatat 8940
aacttaatgg aaataattac ttaaaaggac gacattatta cccatcttcc attaatgata 9000
tattattgat atttaagtta aaccagtaa atgaagtcca tgggaataata gaaagagaaa 9060
aagcattttc aggtataggt gttttgggaa acaatttccc ataataacta taaattcaat 9120
ttgggtcatt tacttcaggt acctatttat ctttctcttt ttcgtaaaag tccatatcca 9180
caaaaccctt tgtaaagggt cgaaccatta tatttctcta catcagaaag gtataaatca 9240
taaaactctt tgaagtcatt ctttacagga gtccaaatac cagagaatgt tttagatata 9300
gcttggtaat ataaagagat gtagtccttc catatttagt attttgagaa acttcagtaa 9360
gaaatgtcct caggtttatg gtctcttaca aaatctatgt ccatcaaaaa ttgtataaag 9420
tggtctaac ttatcccaat aacctaaact tccgtcgcta ttgtaaccag ttctaaaagc 9480
tgtatttgag tttatcaccg ggtagttttt aacatatttc accgagattg aatagggtta 9540
ttggattgag aggcagcgat aacattgggtc aagatttttc acataaactc aaatagtggg 9600
ttgtcactaa gaaaataaat gcagggtaaa atttatatcc ttcttgtttt atgtttcggg 9660
ataaaactt aatatcaatt tctgtggtta tactaaaagt aacagtgatt cttttattta 9720
cgtcccatth taaatatagg aagaacaaaa tacaaagcca tattttgtga ttatagttaa 9780
agacaccaat atgattttca cgttggttggt tcaaaataat gattaaatat ctcttttctc 9840
ttccaattgt ctaaatcaat tttattaaag ttcatttgat atgcctccta aatttttatc 9900
gcaaacaccc aagtttatta ctaatttata gagaaaagag aagggttaaca gatttagtta 9960
aaataatttc aagtaacta tacggaggat ttaaaaaatag taaagtgaat ttaggaggct 10020
tacttgtctg ctttcttcat tagaatcaat ccttttttaa aagtcaatat tactgtaaca 10080
taaatatata ttttaaaaat atttcactta aatcctccga atgaacagac gaaagaagta 10140
atcttagtta ggaaaaaatt tttagttata atgacattgt atttatatat aaaattttta 10200
atcccacttt atccaatatt cgttccttaa tttcatgaac aatcttcatt ctttcttctc 10260
tagtcattat tattggtccc agatctgggt gaactactct taggggtgaaa taggttataa 10320
gcaaggaatt aaagtacttg ttagaagtaa gaaagaagag atcagtaata ataaccaggg 10380
tctagacca cttgatgaga ttaataaaat aatttttccg ttccaattc cacattgcaa 10440
taatagaaaa tccatcttca tcggcttttt cgctcatcat tgtatgaatc aaatcgctt 10500
aattatttta ttaaaaaggc aagggttaag gtgtaacgtt attatctttt aggtagaagt 10560
agccgaaaaa gcagtagtag acatacttag tttagcggaa cttctgtgtc atcaagggtt 10620
aattttttat gtatttcttt taacaaacca ccataggaga ttaacctttt acggtgtaaa 10680
ccttcctcca aatcagacaa gaagacacag tagttccaaa ttaaaaaata cataaagaaa 10740
attgtttggt ggtatcctct aattggaaaa tgccacattt ggaaggagggt ttagtctgtt 10800
acgtttcaaa ttcttttctt catcatcggt cataaaatcc gtatccttta caggatattt 10860

```

```

tgcagtttcg tcaattgccg attgtatatc cgattttatat tgcaaagttt aagaaaagaa 10920
gtagtagcca gtatttttagg cataggaaat gtcctataaa acgtcaaagc agttaacggc 10980
taacatatag gctaaatata ttatttttcg gtcgaatcat ttgaactttt acatttggtat 11040
catagtctaa tttcattgcc tttttccaaa attgaatcca ttgtttttga ttcacgtagt 11100
aataaaaagc cagcttagta aacttgaaaa tgtaaaccta gtatcagatt aaagtaacgg 11160
aaaaagggtt taacttaggt aacaaaaact aagtgcacat tttctgtatt cttaaaataa 11220
gttggttcca cacataccaa tacatgcatg tgctgattat aagaattatc tttattattt 11280
attgtcactt ccgttgccag aaagacataa gaatttttatt caaccaagggt gtgtatgggt 11340
atgtacgtac acgactaata ttcttaatag aaataataaa taacagtga ggcaacgtgc 11400
cataaaacca acaagatttt tattaatttt tttatattgc atcattcggc gaaatccttg 11460
agccatatct gacaaactct tatttaattc ttcgccatca gtattttgggt tgttctaaaa 11520
ataattaaaa aaatataacg tagtaagccg ctttaggaac tcggtataga ctgtttgaga 11580
ataaattaag aagcggtagt taaacatttt taactgttaa tgtgagaaac aaccaacgaa 11640
ctgttggtt ttgtttaata acttcagcaa caaccttttg tgactgaatg ccatgtttca 11700
atgtgtaaaa attgacaatt acactctttg ttggttgctt gacaaccgaa aacaaattat 11760
tgaagtcgtt gttggaaaac actgacttac ggtacaaagt ttgctctcct ccagttgcac 11820
attggacaaa gcctggattt acaaaaccac actcgataca actttctttc gcctgtttca 11880
cgattttggt tatactctaa aacgagagga ggtcaacgtg taacctgttt cggacctaaa 11940
tgttttggtg tgagctatgt tgaaagaaag cggacaaagt gctaaaacaa atatgagatt 12000
tatttcagca caatctttta ctctttcagc ctttttaaata tcaagaatat gcagaagttc 12060
aaagtaatca acattagcga ttttcttttc tctccatggt ataaagtcgt gttagaaaat 12120
gagaaagtcg gaaaaattta agttcttata cgtcttcaag tttcattagt tgtaatcgct 12180
aaaagaaaag agaggtagca ctactttttc cactttttgt cttgtccact aaaacccttg 12240
atttttcatc tgaataaatg ctactattag gacacataat attaaaagaa acccccatct 12300
gagtgaagag gtgaaaaaca gaacagggtg ttttggaac taaaaagtag acttatttac 12360
gatgataatc ctgtgtatta taattttctt tgggggtaga atttagttat ttgtttagtc 12420
acttataact ttaacagatg gggtttttct gtgcaaccaa ttttaagggt tttcaatact 12480
ttaaaacaca tacataccaa taaatcaata aacaaatcag tgaatattga aattgtctac 12540
cccaaaaaga cacgttggtt aaaattccca aaagttatga aattttgtgt atgtatgggt 12600
cacttcaacg cacctttcag caactaaaat aaaaatgacg ttattttctat atgtatcaag 12660
ataagaaaga acaagttcaa aaccatcaaa aaaagacacc gtgaagttgc gtggaaagtc 12720
gttgatttta tttttactgc aataaagata tacatagttc tattctttct tgttcaagtt 12780
ttggtagttt tttctgtgg ttttcagggt ctttttttat tttataaact cattgggtga 12840
tctcgacttc gttctttttt tacctctcgg ttatgagtta gttcaaattc gttcttttta 12900
aaaagtcac gaaaaaata aaatatttga gtaaccact agagctgaag caagaaaaaa 12960
atggagagcc aatactcaat caagtttaag caagaaaaat ggttctaaat cgtgtttttc 13020
ttggaattgt gctgttttat cctttacctt gtctacaaac cccttaaaaa cgttttttaa 13080
ggcttttaag ccgtctgtac ccaagattta gcacaaaaag aaccttaaca cgacaaaata 13140
ggaaatggaa cagatgtttg gggaattttt gcaaaaattt ccgaaaattc ggcagacatg 13200
gttccttaag gcaaggaatt cc
13222

```

<210> 61

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 61

aaggcgccgac acacctacaa catgaggagt agccatcgat gtaccgtaa agctggcata 60
tggtgaac 68

<210> 62

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 62

aaccgcacag cgttttttta ttgattaacg cgttgc 36

<210> 63

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 63

gagttaagcc cagaagatgt ggacgcg 27

<210> 64

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 64

ccgaacctga accatccgcg gccctagga ctttaacagc 40

<210> 65

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 65

catcgatgta ccgtttggtgta agctggcata tgttg 35

<210> 66

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 66

aaccgcacag cgttttttta ttgattaacg cgttgc 36

<210> 67

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 67

gctgttaaag tcctagggat cgcgactggt tcaggttcgg tcagc 45

<210> 68

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 68

gattaacgcg ttgccgcttc tg 22

<210> 69

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 69

gttaaagtc taggggcgtc gagcggttca ggttcggtc

39

<210> 70

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 70

aacgcctcta gatttcgcgc tattaacagc ttgctcgagt gtttcacttg gcgaagggct 60
tcc 63

<210> 71

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 71

ccgaacctga accatccgcg gcccttagga ctttaacagc

40

<210> 72

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 72

gctgttaaag tcctaggggc gggtagcggg tcagggttcgg tc

42

<210> 73

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 73

gattaacgcg ttgccgcttc tg

22

<210> 74

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 74

gtcctcgata cagggatatc cactcatcca gatctaaata ttaaagggtgg cgcaagcttt 60
gta 63

<210> 75

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 75

taggacttta acagc

15

<210> 76

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 76

gctgttaaag tcctaggggc gtcgagcggg tcaggttcgg tcgggtcgat tgcccaagga 60
ttg 63

<210> 77

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 77

gattaacgcg ttgccgcttc tg 22

<210> 78

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 78

gctgttaaag tcctaggggc gtcgggcact ggcagcgggt caggttcggg c 51

<210> 79

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 79

gattaacgcg ttgccgcttc tg 22

<210> 80

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 80

gctgttaaag tcctaggggg cccagccggt tcagggtcgg tcagc 45

<210> 81

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 81

gattaacgcg ttgccgttc tg 22

<210> 82

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 82

gctgttaaag tcctaggggg catccattcg gcagggtcgg tcagctcgat t 51

<210> 83

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 83

gagttaagcc cagaagatgt ggacgcg 27

<210> 84

<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 84
gattaacgcg ttgccgcttc tg 22

<210> 85
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sense primer

<400> 85
gctgttaaag tcctaggggc ggcagacggt tcagggtcgg tcagc 45

<210> 86
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 86
gattaacgcg ttgccgcttc tg 22

<210> 87
<211> 63
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sense primer

<400> 87
gctgttaaag tcctaggggc ggcagacggt tcagggtcgg tcagctcgat tgcccaagga 60
ttg 63

<210> 88

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 88

ttgctcgagt gtggcactgg tcgaagggt tcctaaact

39

<210> 89

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 89

gagttaagcc cagaagatgt ggacgcg

27

<210> 90

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 90

aaccgctcgc ccctgctagg actttaacag

30

<210> 91

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 91
aaccgcacag cgttttttta ttgattaacg cgttgc 36

<210> 92
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sense primer

<400> 92
gagttaagcc cagaagatgt ggacgcg 27

<210> 93
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Antisense
primer

<400> 93
gaccgaacct gaaccgttgc tcgcccctag gac 33

<210> 94
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Sense primer

<400> 94
catcgatgta ccgtttggtg agctggcata tgttg 35

<210> 95
<211> 36
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 95

aaccgcacag cgttttttta ttgattaacg cgttgc

36

<210> 96

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 96

gagttaagcc cagaagatgt ggacgcg

27

<210> 97

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 97

gaccgaacct gaaccatcgc tcgcccctag gac

33

<210> 98

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 98

aaccgcacag cgttttttta ttgattaacg cgttgc

36

<210> 99

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 99

gagttaagcc cagaagatgt ggacgcg

27

<210> 100

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 100

<210> 101

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 101

aaccgcacag cgttttttta ttgattaacg cgttgc

36

<210> 102

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 102

gagttaagcc cagaagatgt ggacgcg

27

<210> 103

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 103

tgtgtaaagt aactcatttg gtgagccag

29

<210> 104

<211> 52

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Antisense
primer

<400> 104

ccgactgcc a ttgcgttcgc atacgacgcc ggggcgctga ttgagcctgc ac

52

<210> 105

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Sense primer

<400> 105

aaccgcacag cgttttttta ttgattaacg cgttgc

36

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 00/00660

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C12N 9/54, C11D 3/386

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C12N, C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 0037621 A1 (NOVO NORDISK A/S), 29 June 2000 (29.06.00) --	1-43
P,X	WO 0037599 A1 (NOVO NORDISK A/S), 29 June 2000 (29.06.00) --	1-43
P,A	WO 0037658 A2 (GENENCOR INTERNATIONAL, INC.), 29 June 2000 (29.06.00) --	1-43
A	US 5691295 A (KARL-HEINZ MAURER ET AL), 25 November 1997 (25.11.97) -----	1-43

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

21 March 2001

Date of mailing of the international search report

06.04.01

Name and mailing address of the International Searching Authority
European Patent Office P.B. 5818 Patentlaan 2
NL-2280 HV Rijswijk
Tel (+31-70)340-2040, Tx 31 651 epo nl,
Fax (+31-70)340-3018

Authorized officer

Carolina Palmcrantz/Eö
Telephone No.

INTERNATIONAL SEARCH REPORT

Information on patent family members

25/02/01

International application No.

PCT/DK 00/00660

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	0037621	A1	29/06/00	AU	1772500 A	12/07/00
WO	0037599	A1	29/06/00	AU	1772800 A	12/07/00
WO	0037658	A2	29/06/00	AU	2200100 A	12/07/00
US	5691295	A	25/11/97	EP	0701605 A	20/03/96
				JP	10507901 T	04/08/98
				US	5855625 A	05/01/99
				WO	9523221 A	31/08/95